

JPG To PDF - Unregistered

If you want to remove this text, Please register

JPG To PDF - Unregistered

If you want to remove this text, Please register

JPG To PDF - Unregistered

If you want to remove this text, Please register

JPG To PDF - Unregistered

If you want to remove this text, Please register



Original research article

Investigating the nonlinear behavior of cobalt (II) phthalocyanine using visible CW laser beam

Ayat J. Kadhum^a, Nazar A. Hussein^a, Qusay M.A. Hassan^{b,*}, H.A. Sultan^b,
Ahmed S. Al-Asadi^b, C.A. Emsbahy^b

^a Department of Chemistry, College of Education for Pure Sciences, University of Basrah, Basrah, Iraq
^b Department of Physics, College of Education for Pure Sciences, University of Basrah, Basrah, Iraq



ARTICLE INFO

Article history:
Received 12 October 2017
Accepted 20 November 2017

MSC:
42.70.-4
42.05.-4
42.05.Aa

Keywords:
Phthalocyanine
Self-phase modulation
Thermal nonlinearity
Diffraction ring pattern
Z-scan technique

ABSTRACT

Experimental and theoretical results of the diffraction ring patterns in cobalt (II) phthalocyanine (PVCuCl₂) in dimethylformamide solution using visible low power continuous wave CW laser beam are reported. The wave-front curvature of the used laser beam seem to modify the spatial phase modulation of the beam. The upward-convection modify each pattern by reducing the vertical diameter of each ring in comparison with horizontal size. As low by reducing the effect of convection is minimized. Simulation results given are based on the Fresnel-Kirchhoff diffraction integral. The nonlinear refractive index of the solution of PVCuCl₂ was measured using Z-scan technique, by exciting with CW laser at 473 nm wavelength. The effect of concentration of the sample on nonlinear refractive index has been investigated.

© 2017 Published by Elsevier GmbH.

1. Introduction

Organic molecules with nonlinear optical process continue to attract attention because of their potential applications in a wide variety of optoelectronic and photonic devices [1–3]. The nonlinear response of these organic molecules at low intensity lasers results in a large third order nonlinear optical (NLO) susceptibilities. Extensive work has been carried out and the reports have been published for the organic molecules in liquid solutions [4,5].

When a laser beam propagates in an absorbing medium a spatial distribution of temperature is generated according to the light beam irradiance spatial profile. Since the local refraction index thermal lens (TL) [6] is induced. As a direct consequence of the TL effect and the change of phase of the propagating laser beam diffracting ring patterns (DRPs) [7] resulted. Based on the (TL) effect the nonlinear refractive index, n_2 , of the nonlinear medium can be evaluated. The Z-scan technique, developed by Sheik Bahae et al. [8] is another simple and effective tool that can be used to measure the nonlinear refractive index, n_2 .

During the last fifteen years, extensive works have been directed towards the study of nonlinear properties of phthalocyanine, viz., bis-phthalocyanine [9], alkyl phthalocyanine [10], Cu(II) phthalocyanine [11], alkyl phthalocyanine [12], 4-tetra and oct-substituted lead phthalocyanine [13], copper phthalocyanine [14], zinc phthalocyanine [15–17], Lanthanide

* Corresponding author.
E-mail address: qusaym@uobasrah.edu.iq (Q.M.A. Hassan).