

## Q (?

### Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy Volume 223, 5 December 2019, 117297

# Formation and temporal evolution of diffraction ring patterns in a newly prepared dihydropyridone

Rita S. Elias <sup>a</sup>, Qusay M.A. Hassan <sup>b</sup>  $\stackrel{ riangle}{\sim}$   $\stackrel{ riangle}{\sim}$ , C.A. Emshary <sup>b</sup>, H.A. Sultan <sup>b</sup>, Bahjat A. Saeed <sup>c</sup>

#### **⊞ Show more**

https://doi.org/10.1016/j.saa.2019.117297

Get rights and content

## Highlights

- A dihydropyridone has been prepared from butylamine and curcumin.
- The nonlinear optical properties of dihydropyridone are studied.
- Nonlinear refractive index of sample is determined using diffraction ring patterns and Z-scan techniques.
- Sample exhibits self-diffraction ring patterns due self-phase modulation.

#### Abstract

A dihydropyridone has been prepared from butylamine and curcumin. A theoretical DFT study was conducted to determine the most stable conformer of the studied molecule (among three conformers) using the B3LYP/6-311+G(d,p) level of theory. This is assisted by the prediction of the <sup>13</sup>C NMR chemical shifts of the conformers which then correlated with the observed <sup>13</sup>C NMR chemical shifts. A TD-DFT study was conducted to analyze the electronic spectrum of the most stable conformer in order to determine the transitions responsible for the longer band in the electronic spectrum of the molecule. As well the frontier orbitals in the most stable conformer were analyzed to establish the density of donor and acceptor sites in the molecule that may be responsible for the nonlinear optical (NLO) properties of the studied molecule.

**no** de la beam quartersed shaffin cell containing solution of dihydropyridone. The nonlinear refractive index,  $n_2$ , was determined based on the number of diffraction rings per a pattern observed and by the Z-scan technique and both results are compared. The upward convection heat effect appears to be responsible for the asymmetries observed in the diffraction ring patterns. The use of convergent and divergent laser beams has led to new types of diffraction ring patterns. Temporal evolution of each diffraction ring patterns was registered. The diffraction ring patterns experimentally obtained are numerically calculated using the Fresnel-Kirchhoff diffraction integral, with good qualitative and reasonable quantitative agreements.

#### Graphical abstract





Next >

#### Keywords

Self-phase modulation; Thermal nonlinearity; Nonlinear refractive index; Diffraction ring pattern; Z-scan technique

Recommended articles Citing articles (0)

© 2019 Published by Elsevier B.V.

ELSEVIER About ScienceDirect Remote access Shopping cart Advertise Contact and support Terms and conditions Privacy policy :=

#### GetyAucesse to the have of cookies ort

Copyright © 2019 Elsevier B.V. or its licensors or contributors. ScienceDirect ® is a registered trademark of Elsevier B.V.

