

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/320730050>

EFFECT OF LIGHT COLOR AND STOCKING DENSITY ON INTESTINAL MORPHOLOGY OF BROILER CHICKENS

Article · March 2017

CITATIONS

0

READS

100

3 authors, including:



Muhsin Sagheer

University of Basrah

11 PUBLICATIONS 0 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



effects of light color and stocking density on intestinal morphological [View project](#)



ISSN: 0976-3376

Available Online at <http://www.journalajst.com>

ASIAN JOURNAL OF
SCIENCE AND TECHNOLOGY

Asian Journal of Science and Technology
Vol. 08, Issue, 03, pp.4340-4344, March, 2017

RESEARCH ARTICLE

EFFECT OF LIGHT COLOR AND STOCKING DENSITY ON INTESTINAL MORPHOLOGY OF BROILER CHICKENS

^{1,*}Mudhar A. S. Abu Tabeeekh, ²Muhsin S. G. Al-Moziel and ³Huda F. Saad

¹Basra Veterinary Hospital, Basra, Iraq

²Department of Physiology and Pharmacology, College of Pharmacy, University of Basra

³Department of Animal Resources, College of Agriculture, University of Basra

ARTICLE INFO

Article History:

Received 18th December, 2016

Received in revised form

27th January, 2017

Accepted 20th February, 2017

Published online 31st March, 2017

Key words:

Broilers, Light color,
Small intestine, Stocking density.

ABSTRACT

Many studies have been conducted to determine the effect of light on broiler performance, and many others on the effect of stocking density. However, there has been a very few investigation on the influence of light color on intestinal morphology of broilers. This study intended to evaluate the effects of color lights and bird density as it relates to small intestine (villi length and crypt depth) of broilers. A total of 675 Ross 308 one-day-old broiler chicks were used in this study. The birds were exposed to white light (WL) as a control, red light (RL), blue light (BL), green light (GL), and Blue – Green mix light (BGL) by a light-emitting diode system (LED) applied for 24 hours daily in separated rooms. The birds were randomly divided and housed into 9 wooden sealed pens of 1m² in three replicates for each density 12, 15 and 18 birds/m² in the room.

Copyright©2017, Mudhar Abu Tabeeekh et al., This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

INTRODUCTION

Many studies have been shown the effects of light and stocking density on broilers performance in experimental assay, but we decided to evaluate the effects of different color lights and densities on intestinal morphology. The small intestine is the main site of feed digestion, nutrients absorption and secretion of digestive fluids (Yamauchi, 2002). The mucus layer varies in thickness and composition throughout the gut and protects the underlying enterocytes lining the villus from mechanical, enzymatic, and chemical challenges in addition to being a source of intestinal lubrication (Sharma and Schumacher, 1995). Long villi are usually equated with excellent gut health, high absorptive efficiency and healthier intestinal tract of chickens (Alfaro et al., 2007). Under magnification, intestinal villi and crypts are seen. In general, intestinal villi are 0.5-1.5 mm long and they are projecting out of the lamina and epithelium into the lumen of the small intestine. The base of the intestinal villi is covered by young epithelial cells which are known as intestinal crypts (Yamauchi, 2002). Lighting is a powerful exogenous factor in control of many physiological and behavioral processes (Leven, 2000). Light may be the most critical of all environmental factors to birds (Manser, 1996) and consist of three different aspects: quantitative aspect (photoperiod) as

well as qualitative aspects include lighting color (wavelength) and light intensity (Heshmatollah, 2007). Day light has relatively wavelengths between 400 and 700 nm. Birds sense light through their eyes (retinal photoreceptors) and through photosensitive cells in the brain (extra-retinal photoreceptors) (El-Fiky et al., 2008). The associated colors are Blue B (435-500 nm), Green G (500-565 nm) and Yellow Y (500-600nm), Orange O (600-630 nm) and Red R (630-700 nm) (Hakan and Ali, 2005). Affection of light on activity and reproduction of chickens had been known from many years ago and it was also shown by many researcher. A few studies have explored on color light preferences of broiler chicks. Taylor et al. (1969) observed that one day-old chicks prefer red and yellow lights in comparison with blue light. Rozenboim et al. (2004) referred that broilers reared under blue or green light were significantly heavier than those reared under red or white light. Heshmatollah, (2007) reported when chickens had the ability to choose among red, orange, yellow, or green lights, they spent significantly more time under green light. Their second preference was yellow light. Cao et al. (2008) described that, monochromatic light could significantly affect broiler growth and development. The GL promoted broiler growth better at early stage, and the BL promoted broiler growth better at a later stage under an illumination intensity of 15 lx. Stocking density is a much discussed topic in animal science. Increasing stocking density generally leads to a decrease in welfare in many farm animal species (Petherick and Phillips, 2009). Stocking density is calculated by different ways, sometimes

*Corresponding author: Mudhar A. S. Abu Tabeeekh,
Basra Veterinary Hospital, Basra, Iraq.

stocking density is reported using the number of birds per unit area or the amount of area per bird. Currently many companies calculate stocking density by the pound. Instead of being expressed as the number of birds per unit area, density is calculated as bird weight per unit area (Fairchild, 2005). In broilers, high densities have been associated with a decline in body weight (BW), feed intake (FI) and feed conversion rate (FCR) (Sanotraet *et al.*, 2001).

MATERIALS AND METHODS

Animals and animal husbandry

The research was done at the poultry farm, College of Veterinary Medicine, Basra University, Basra, Iraq. A total of 675 Ross 308 one-day-old broiler chicks were used in this study. The chicks were raised under control condition from day one until 35 days of age. Broiler chicks were reared into five light groups in separated rooms 3 x 3 x 4 meters with an average 135 chicks in each room under LED color lights include : White light as a control, Red light (660 nm), Blue light (480 nm), Green light (560 nm) and Blue – Green mix light. Stocking density (12, 15 and 18 birds/m²) were randomly housed into 9 wooden sealed pens of 1m² in three replicates in the room. Light sources were equalized on the intensity of 5 watt/ m² (20 lux) at bird head level and light period of 24 hours daily. Room temperature was initially 34°C which was subsequently reduced by 2°C/week to 22°C by 35 day. Three- dietary pellet rations were used to feed the chicks, which consist of starter, grower, and finisher diets. Total dietary metabolic energy for the starter, grower and finisher were 2925, 3111 and 3171 kcal/kg respectively, while the values of crude protein were 22.21, 20.14 and 18.08 % respectively. Half cylinder plastic feeders were placed in each pen. The birds were supplied with feed and water ad libitum, and diets were formulated to meet the nutrient recommendations for poultry according to NRC, (1994). Nipple water drinking system was set up in each pen and was manually adjusted as birds grew to ensure the watering system was kept at a proper level.

Intestinal mucosal morphology

On days 35, three chickens were randomly selected from each replicate and slaughtered through a section in the jugular vein. The carcass was eviscerated manually by cutting around the vent to remove all of the viscera including the small intestine (Toghyani *et al.*, 2012). Pieces of 1 cm small intestine specimens were fixed in 10% buffered formalin. Each piece of tissue was trimmed at the thickness of 5 microns (μ) in size, fixed and dehydrated in a series of increasing alcohol concentration, and embedded in paraffin wax using rotary tissue processor. The sections were stained with Mayer's Hematoxylin and eosin for observation with a light microscope (Luna, 1968). Villus height was measured as the length between the tip of the villus and the villous-crypt axis. Measurements for crypt depth were taken from the valley between individual villi to the baso- lateral membrane as described by Baurhoo *et al.* (2007). Values are means from 10 different villi and only vertically oriented villi and crypts were measured (Uni *et al.*, 2003) and the ratio of villus length to crypt depth (V:C ratio) was calculated.

Models of Analysis

Data was analyzed using completely randomized design (CRD) according to SPSS, (2009). The significant tests for the differences between each two means for any studied trait were done according to Duncan's multiple rang test. The model was:

$$Y_{ijk} = M + Li + Dj + e_{ijk}$$

Where: Y_{ijk} = Observation on the ij individual

M = Overall mean

Li = light effect

Dj = density effect

e_{ijk} = Random error

RESULTS AND DISCUSSION

Over the last decades, there has been increased interest in intestinal growth/health as poultry nutritionists have attempted to adopt new approaches to deal with the broader changes in the overall agricultural landscape. Color is an important aspect of light that has been considered at one time as a management tool in poultry production (Prayitno, 1997). The results of this study revealed a significant increase ($P < 0.05$) of villus height 786 μm in broilers reared under WL but no significant effect ($P > 0.05$) was recorded under different levels of stocking density as shown in Table 1. The significant effect of WL on feed intake was confirmed by Rierison, (2011) who found that birds during weeks 1 to 3, prefer white light. The chicks demonstrated a preference for pelleted feed and white light ($P < 0.01$) and they chose not to feed under green or blue light. A possible explanation as to why broilers prefer to consume feed under white light could be due to it helps them to identify texture differences they cannot see under different colors. Adopting a strategy allowing broiler chicks to feed under white light and rest under blue or green light. Prayitno *et al.* (1997) conducted a study to compare broilers behavior under red, white, blue, or green lights. It was concluded that broilers raised under red and white lights were more active than those raised under blue and green lights as expressed by greater walking activity in the white light treatment. It has been suggested that feed intake outstrips the increase in size of the gastro intestinal tract (GIT) and therefore rate of passage of feed through the intestine is faster (Sibbald, 1979). Ferket and Gernat (2003); Forbes (2005); Scott, (2005) agreed that voluntary feed intake is linked to growth rate. Two excellent reviews on the biological relationship between intestinal development and growth in avian species can be found in Konarzewski *et al.* (1989; 1990). Sklan *et al.* (1975) ; Sklan, and HuIwitz, (1980) referred to the increases in length, internal diameter, and villus volume of the small intestine together with the five to eight-fold increase in feed. Overall, the increased feed intake with age is accompanied by rapid development of the GIT, especially in the first week after hatch (Noy and Sklan, 1997). Karakaya *et al.* (2009) found that, feed consumption, body weight and total muscle weight values of the muscles from GL-BL and GL-GBL mix lighting groups were significantly higher than those of incandescent (control) lighting groups. Cao *et al.* (2008) indicated that broilers reared under BL were larger in the weights of carcass, breast muscle, thigh, crus, and net chamber at d 49 as compared with other light groups.

Table 1. Effect of light color and stocking density on Villus height (μm) of broiler intestine at 35 day ($M \pm SE$)

Light color	WL	RL	BL	GL	BGL	Effect of stocking density
Stocking density						
12 birds/m ²	^{ab} 685.8 $\pm$ 17.63	^a 777.6 $\pm$ 15.74	^a 763.2 $\pm$ 42.53	^b 651.6 $\pm$ 25.55	^c 473.4 $\pm$ 11.24	670.3 $\pm$ 24.51
15 birds/m ²	^a 837.0 $\pm$ 24.48	^b 565.2 $\pm$ 32.37	^a 723.6 $\pm$ 46.47	^a 819.0 $\pm$ 9.85	^a 808.2 $\pm$ 15.16	750.6 $\pm$ 23.64
18 birds/m ²	^a 835.2 $\pm$ 7.74	^b 626.4 $\pm$ 30.99	^b 572.4 $\pm$ 34.22	^a 754.8 $\pm$ 12.67	^a 743.4 $\pm$ 31.64	706.44 $\pm$ 21.99
Effect of light color	^a 786.0 $\pm$ 21.23	^b 656.4 $\pm$ 28.00	^{ab} 686.4 $\pm$ 31.18	^{ab} 741.8 $\pm$ 20.64	^{ab} 675.0 $\pm$ 40.38	N.S.

*a, b, c Means in horizontal and vertical rows with different superscripts were significantly different at ($p < 0.05$). * N.S. not significant. SE: standard error

Table 2. Effect of light color and stocking density on Crypt depth (μm) of broiler intestine at 35 day ($M \pm SE$)

Light color	WL	RL	BL	GL	BGL	Effect of stocking density
Stocking density						
12 birds/m ²	^{ab} 147.6 $\pm$ 14.4	^a 172.8 $\pm$ 10.41	^a 176.4 $\pm$ 12.91	^{ab} 138.6 $\pm$ 8.34	^b 106.2 $\pm$ 3.36	^b 148.32 $\pm$ 6.77
15 birds/m ²	^a 217.8 $\pm$ 8.72	^b 131.4 $\pm$ 8.34	^a 210.6 $\pm$ 10.10	^a 205.2 $\pm$ 7.20	^a 203.4 $\pm$ 9.26	^a 193.68 $\pm$ 7.36
18 birds/m ²	^a 214.2 $\pm$ 6.61	^{bc} 158.4 $\pm$ 8.34	^c 147.6 $\pm$ 14.40	^{ab} 194.4 $\pm$ 8.34	^{ab} 187.2 $\pm$ 5.24	^a 180.36 $\pm$ 6.20
Effect of light color	193.2 $\pm$ 10.27	154.2 $\pm$ 6.68	178.2 $\pm$ 9.62	179.4 $\pm$ 8.89	165.6 $\pm$ 11.87	*

*a, b, c Means in horizontal and vertical rows with different superscripts were significantly different at ($p < 0.05$). * N.S. not significant. SE: standard error

Table 3. Effect of light color and stocking density on V:C ratio of broiler intestine at 35 day ($M \pm SE$)

Light color	WL	RL	BL	GL	BGL	Effect of stocking density
Stocking density						
12 birds/m ²	4.01 $\pm$ 0.88	4.54 $\pm$ 0.20	4.34 $\pm$ 0.12	4.75 $\pm$ 0.27	4.47 $\pm$ 0.18	^a 4.58 $\pm$ 0.11
15 birds/m ²	^{ab} 3.85 $\pm$ 0.09	^a 4.31 $\pm$ 0.16	^b 3.43 $\pm$ 0.16	^{ab} 4.00 $\pm$ 0.12	^{ab} 3.99 $\pm$ 0.13	^b 3.92 $\pm$ 0.08
18 birds/m ²	3.90 $\pm$	3.96 $\pm$	3.94 $\pm$ 0.23	3.90 $\pm$ 0.12	3.97 $\pm$ 0.13	^b 3.93 $\pm$ 0.06
Effect of light color	0.09	0.12	3.91 $\pm$ 0.13	4.21 $\pm$ 0.14	4.14 $\pm$ 0.10	*

*a, b, c Means in horizontal and vertical rows with different superscripts were significantly different at ($p < 0.05$). * N.S. not significant. SE: standard error

According to Xie *et al.* (2009), who studied Arbor Acres male broilers were exposed to red, green, blue and white light from LED lamps. The study found that small intestine mucosal structure is improved to an extent in broilers when illuminated with green light at the early growth stage (0-7 days) under 15 Lux light intensity, accordingly improving the absorption function of the small intestine and accelerating growth. Xie *et al.* (2011), found that villus height of small intestine was increased in broilers exposed to GL and BL than those in the WL group. The suggestion that both mucosal mechanical and immunological barriers of the small intestine might be improved by rearing broilers under GL at an early age and under BL at an older age. The results of Firouzi *et al.* (2014) indicated that the birds reared under yellow and blue light had the best and weakest performance, respectively. According to results in Table 2 and 3, no significant difference ($P > 0.05$) of color light was observed between treatments considering the crypt depth (μm) and V:C ratio of broiler intestine, while a significant effect ($P < 0.05$) of treatments was detected for the stocking density which recorded 193.68 μm in the treatment of 15 birds/m² for crypt depth and 4.58 in the treatment of 12 birds/m² for V:C ratio. Increased stocking densities are frequently reported to depress chicken growth performance, but the mechanisms behind this are not fully understood (Guardia, 2011). Riyad *et al.* (2014) revealed that The small intestine length (cm) was differed ($P < 0.05$) in broilers reared under GL and 18 birds/m², while small intestine weight (gm) was higher ($P < 0.05$) under BL and 12 birds/m² in compare with other groups at 35th day. Alaeldein *et al.* (2013) revealed that heavier breasts were obtained from birds which had subjected to the low stocking density ($p < 0.001$). Total small intestine lengths and weights from birds which were subjected to the low density were the longest and the heaviest as compared to the other two groups.

Burkholder *et al.* (2008) reported that stressful conditions had a negative impact on intestinal morphology. A negative effect (Dozier *et al.*, 2006), or a positive effect in the first 3 week (Moreira *et al.*, 2004). These discrepancies between studies concerning the effect of stocking density on growth performance of the chicken during the first 3 week of life may be explained by the different experimental conditions, such as bird strain, the presence or absence of antibiotics, or litter type (Guardia, 2011).

Conclusion

Based on the results of the present study, it can be stated that light color light and stocking density affected broiler intestinal morphology. However, results from previous studies are not completely consistent, especially with respect to the use of colored light. These conflicts were probably due to differences of light source, light schedule, animal species, and age of experimental animals used in different investigations as well as different experimental conditions of stocking density.

REFERENCES

- Alaeldein M. Abudabos, Mutassim M. Abdelrahman, Hany M. Yehia, Muath Q. Al-Ghadi and Ibrahim A. Alhidary, 2013. Effect of Stocking Density on Intestinal Histology and Ileal Bacterial Count in Broilers. *Asian Journal of Animal and Veterinary Advances*, 8: 740-746.
- Alfaro, D.M., Silva, A.V.F., Borges, S.A., Maiorka, F.A., Vargas, S. and Santin, E. 2007. Use of *Yucca schidigera* extract in broiler diets and its effects on performance results obtained with different coccidiosis control methods. *J. Applied Poult. Res.*, 16: 248-254.

- Baurhoo, B., Phillip, L. and Ruiz-Feria, C.A. 2007. Effects of purified lignin and mannan oligosaccharides on intestinal integrity and microbial populations in the ceca and litter of broiler chickens. *Poultry Science*, 86: 1070–1078.
- Burkholder, K.M., Thompson, K.L., Einstein, M.E., Applegate, T.J., Patterson, J.A. 2008. Influence of stressors on normal intestinal microbiota, intestinal morphology, and susceptibility to *Salmonella enteritidis* colonization in broilers. *Poultry Science*, 87: 1734–1741.
- Cao, J. , Liu, W. , Wang, Z. , Xie, D. and Chen, Y. 2008. Green and blue monochromatic lights promote growth and development of broilers via stimulating testosterone secretion and microfiber growth. *J. Appl. Poult. Res.*, (17):211–218.
- Dozier, W.A., Thaxton, J., Purswell, J.L., Olanrewaju, H.A., Branton, S. and Roush, W. 2006. Stocking density effects on male broilers grown to 1.8 kilograms of body weight. *Poult. Sci.*, 85:344–351.
- El-Fiky, A., Soltan, M., Kalamah, M. A. and Abou-Saad, S. 2008. Effect of light color on some productive, reproductive, egg quality traits, and free radicals in turkey. *Egypt. Poult. Sci.*, Vol 28 (3): 677–699.
- Fairchild, B. D. 2005. Broiler stocking density, Cooperative Extension Service College of Agricultural and Environmental Sciences, The University of Georgia, USA.
- Ferket, P.R. and Gernat, A.G. 2003. Feed intake A–Z. *Watt Poultry USA*, pp. 14–29.
- Firouzi, S, Nazarpak H.H, Habibi, H., Jalali, S.S., Nabizadeh, Y., Rezaee, F., Ardali, R., Marzban, M. 2014. Effects of Color Lights on Performance, Immune Response and Hematological Indices of Broilers. *J. World's Poult. Res.* 4(2): 52–55.
- Forbes, J.M. 2005. Why did the chicken choose the food? Australian Poultry Science Symposium 17, 145–152.
- Guardia, S., Konsak, B., Combes, S., Levenez, F., Cauquil, L., Guillot, J.-F., Moreau-Vauzelle, C., Lessire, M., Juin, H., and Gabriel, I. 2011. Effects of stocking density on the growth performance and digestive microbiota of broiler chickens. *Poultry Science*. 90 (9):1878–1889.
- Hakan, B. and Ali, A. 2005. Effects of light wavelength on broiler performance. *Hayvansal Uretim*. (46): 22–32.
- Heshmatollah, K. 2007. Preference of Broiler Chicks for Color of Lighting and Feed. *Journal of Poultry Science*, 44: 213–219.
- Karakaya, M., Parlatb, S.S., Yilmaza, M.T., Yildirim, I. and Ozalpa, B. 2009. Growth performance and quality properties of meat from broiler chickens reared under different monochromatic light sources. *Brit. Poult. Sci.*, 50:76–82.
- Konarzewski, M., Kozłowski, J. and Zioiko, M. 1989. Optimal allocation of energy to growth of alimentary tract in birds. *Funct. Ecol.*, 3:589–596.
- Konarzewski, M., Lilja, C., Kozłowski, J. and Lewonczuk, B. 1990. On the optimal growth of the alimentary tract in avian postembryonic development. *J. Zool. Lond.*, 222:89–101.
- Leven, M. N. 2000. The Cosmic Rainbow, Second edition, Health Arena Publishing and Promotions, 19 Butler Street, Willagee, Western Australia. Pp 1–7.
- Luna, L.G. 1968. Manual of histologic staining methods of the Armed Forces Institute of Pathology. 3rd Ed. New York: Mac Graw-will. 258p.
- Manser, C.E. 1996. Effects of lighting on the welfare of domestic poultry: a rev. *Anim. Welf.*, (5):341–360.
- Moreira, J., Mendes, A. A., Roca, R. O., Garcia, E. A., Naas, I. A., Garcia, R. G., Paz, I. C. L. A. 2004. Effect of stocking density on the performance, carcass yield and meat quality of broilers of different commercial strains. *R. Bras. Zootec.*, 33:1506–1519.
- Noy, Y. and Sklan, D. 1997. Posthatch Development in Poultry. *J. Appl. Poult. Res.*, 6: 344–354.
- NRC, 1994. Nutrient Requirements of Poultry. 9th rev. ed. Natl. Acad. Press, Washington DC, USA.
- Petherick, J.C. and Phillips, C.J.C. 2009. Space allowances for confined livestock and their determination from allometric principles. *Appl. Anim. Behav. Sci.*, (117): 1–12.
- Prayitno, D. S.; Phillips, C.J.C. and Omed, H. 1997. The Effects of Color of Lighting on the Behavior and Production of Meat Chickens. *Poult. Sci.*, (76):452–457.
- Rierson, R. D. 2011. Broiler performance for light color and feed form, and the effect of light on growth and performance of broiler chicks. Master of Science, College of Agriculture, Kansas State University, Manhattan, Kansas, USA.
- Rozenboim, I., Biran, I., Chaiseha, Y., Yahav, S., Rosenstrauch, A., Sklan, D., Halevy, O. 2004. The Effect of a Green and Blue Monochromatic Light Combination on Broiler Growth and Development. *Poultry Science*, 83:842–845.
- Riyad K. Mosa, Rabia J. Abbas, Mudhar A. S. Abu Tabeekh. 2014. The effect of color light and stocking density on some traits of broiler carcasses. *MRSVA*. 3 (2), 24–35.
- Sanotra, G. S., Lawson, L. G. and Vestergaard, K. S. 2001. Influence of stocking density on tonic immobility, lameness, and tibial dischondroplasia in broilers. *J. Appl. Anim. Welf. Sci.*, (4):71–87.
- Scott, T.A. 2005. Variation in feed intake of broiler chickens. *Recent Advances in Animal Nutrition in Australia*, Volume 15. 237–244.
- Sharma, R. and Schumacher, U. 1995. Morphometric analysis of intestinal mucins under different dietary conditions and gut flora in rats. *Dig. Dis. Sci.*, 40:2532–2539.
- Sibbald, I., Jr., 1979. Passage of feed through the adult rooster. *Poultry Sci.*, 58:446–459.
- Sklan, D., Hudtz, S., Budowski, P. and Ascarelli, I. 1975. Fat digestion and absorption in chicks fed raw or heated soybean meal. *J. Nutr.*, 105:5743.
- Sklan, D. and Hulwitz, S. 1980. Protein digestion and absorption in young chicks and turkeys. *J. Nutr.*, 110:139–144.
- SPSS, 2009. Statistical Package of Soc. Sci., Ver.18. Appl. Guide. Copy right by SPSS Inc. USA.
- Taylor, A., Sluckin, W. and Hewitt, R. 1969. Changing color preferences of chicks. *Animal Behavior*, 17: 3–8.
- Toghyani, M., Tohidi, M., Gheisari, A., Tabeidian, A. and Toghyani, M. 2012. Evaluation of Oyster Mushroom (*Pleurotus ostreatus*) as a Biological Growth Promoter on Performance, Humeral Immunity, and Blood Characteristics of Broiler Chicks. *J. Poult. Sci.* 49: 183–190.
- Uni, Z. Smirnov, A. and Sklan, D. 2003. Pre-and posthatch development of goblet cells in the broiler small intestine; Effect of delayed access to feed. *Poult. Sci.*, 82(2):320–327.
- Xie, D, Chen, Y.X. , Wang, Z. X. and Dong, Y.L. 2009. Effects of monochromatic light on structure of small intestinal mucosa in broilers. *Scientia Agricultura Sinica* 42 (3): 1084–1090.

- Xie, D., Li, J., Wang, Z. X., Cao, J., Li, T. T., Chen, J. L. and Chen, Y.X. 2011. Effects of monochromatic light on mucosal mechanical and immunological barriers in the small intestine of broilers. *Poult. Sci.*, 90 (12): 2697-2704.
- Yamauchi, K. 2002 Review on chicken intestinal villus histological alterations related with intestinal function. *Journal of Poultry Science*, 39: 229-242.
