



ISSN 2277-386X

Original Article

Estimation of Polycyclic Aromatic Hydrocarbons (PAHs) in Some Fishes of Shatt Al-Arab River-Iraq

Hamid T. Al-Saad*Dhafar Dh. AL-Khion and Balqees S. Al-Ali

Department of Marine environmental Chemistry, Marine Science Center, University of Basrah, Basrah, Iraq

*Corresponding Author= htalsaad@yahoo.com

Received 17 July 2016; accepted 23 August 2016

Abstract

Levels of sixteen polycyclic aromatic hydrocarbons (PAHs) in six edible tissues of selected frequently-consumed fish collected from Shatt Al-Arab river, namely Cattana (*Luciobarbus xanthopterus*), Gareeba (*Ctenopharyngodon delta*), Samti (*Cyprinus carpio*), Bultti (*Tilapia zillii*), Khishni (*Palaniza abu*) and Shillig (*Leuciscus vorax*) were investigated in 2015. High Pressure Liquid Chromatography (HPLC) analysis was employed for determination of PAHs. Total PAHs in the samples ranged from 0.02ng/g dry weights in *Cy. carpio* to 3.34ng/g dry weights in *Le. vorax*. Some ratios were calculated to evaluate the sources of these compounds in the area. The results showed that there was pyrolytic and petrogenic sources. Low molecular weight (LMW) PAHs was the predominant compounds, suggesting the gill-water transfer might be the major exposure route for PAHs in the studied fish species.

© 2016 Universal Research Publications. All rights reserved

Keywords:- Shatt Al-Arab River, Polycyclic aromatic hydrocarbons (PAH), Edible fishes, sources.

Introduction

Water is an essential natural resource for sustaining the environment and life, that cycle through various reservoirs within and among ecosystems. Thus any effect upon resource due to addition of some materials and can be harmful to living organisms referred to pollution[1]. Water resources play a vital role in various sectors of economy such as agriculture, livestock production, industrial activities, hydropower generation, fisheries and other creative activities. It has been put under tremendous pressure, and has been deteriorated due to some important factors like increasing urbanization, industrialization, and agriculture activities [2].

The Shatt Al-Arab River is formed after the confluence of the Euphrates and the Tigris Rivers near the city of Qurna in southern Iraq. The southern part of the river constitutes the border between Iran and Iraq until it discharges into the Arabian Gulf, with a total length of 192 km, the Shatt Al-Arab widens over its course, expanding from a width of 250-300 m near the Euphrates-Tigris confluence to almost 700 m near the city of Basrah and more than 800 m as it approaches the river mouth [3]. The Shatt al Arab Delta is 140 km wide and splits into more than 10 branches. The landscape is characterized by green marshy areas, lakes, lagoons and estuaries, bordered by irrigated lands and date

palm plantations and surrounded by desert[4].

Fish exposed as marine biologist at risk of oil pollution, which may be fatal as that of oil products such as gasoline or diesel oil or kerosene, causing mass death of fish and other aquatic organisms [5].

Polycyclic Aromatic Hydrocarbons (PAHs) is another large group of hydrocarbons materials with two or more fused aromatic rings. PAHs contamination is present in incomplete combustion of organic and fossil fuel as phenanthrene, fluorine, benzo[a]pyrene and pyrene in car exhaust, while phenanthrene, fluorine, and pyrene in diesel vehicles exhaust [6]. Naphthalene and acenaphthalene have been reported to be present in fuel oil used extensively in Shatt Al-Arab region, and these compounds are not easily attacked by microorganisms, and numerous of these compounds as well as their derivatives are potential carcinogens[7]. The solubility of these compounds in the water is low and they tend to be bounded to suspended organic matter in the water column and finally accumulate in the marine sediment[8].

Due to the large number of studies on aromatic compounds in marine fish [9] so that it is necessary to study these compounds in fresh fish because of their economic importance. Therefore, the aim of the present study was to evaluate and determine 16 compounds of PAHs residues in

six fresh commercial fish species at Shatt al-Arab River.

Materials and methods

Six species of fish samples which were commonly available at Shatt Al-Arab River, namely Cattar (*Luciobarbus xanthopterus*), Gareeba (*Ctenopharyngodon idella*), Samti (*Cyprinus carpio*), Bultti (*Tilapia zillii*), Khishni (*Palanizaabu*) and Shillig (*Leuciscus vorax*) were collected during 2015 by the local fishermen using drag net (Mesh size 1.27 cm, thickness 9 ply). The samples were kept refrigerated in polyethylene boxes with ice during transport to the laboratory. The fish samples then stored at -20°C until further analysis. Samples were removed from the deep freezer, thawed, and cleaned very well in tap water to remove any dirt. Dissection was performed on thawed fish using aseptic instrument and glass dishes, muscle tissues were dissected. Each fish sample was cut into pieces and crushed in a pestle mortar. The procedure of [10] was used for the extraction of hydrocarbon compounds from fish muscles. The concentrations of PAHs in muscles were determined by using High Pressure Liquid Chromatography (HPLC) Shimadzu CBM20A with UV-visible detector. The fish samples were analyzed for a suite of 16 PAHs; naphthalene (NAP), acenaphthylene (Acy), acenaphthene (Ace), Fluorene (Flu), Phenanthrene (Phe), anthracene (Ant), Fluoranthene (Flut), pyrene (Pyr), Benzo[a]anthracene (BaA), chrysene (Chr), benzo(b)fluoranthene (BbF), Benzo[k]fluoranthene (BkF), benzo(a)pyrene (BaP), Dibenzo[a,h]anthracene (DahA), benzo(ghi)perylene (BPY) and Indeno[1,2,3-cd] pyrene (InP). Some of ratios used to determine sources of PAHs compounds which including the following: (Flut / Pyr), (Phe / Ant), (Ant / (Ant + Phe)), (BaP / (BaP + Chr)) and (LMW / HMW). The calculation of these concentrations can be an indicator of the source of the aromatic hydrocarbons found in the fish samples, ratio of (Flut / Pyr) Values greater than 1 have been used to indicate pyrogenic origins and values less than 1 are attributed to petrogenic source [11]. Ratio of (Phen/Ant) less than 1 indicates petrogenic [12]. ratio of (LMW / HMW) values greater than 1 indicate petrogenic origins and values less than 1 are attributed to pyrogenic sources [13].

(Ant/ (Ant+Phe)) ratio presumes that ratios less than 0.1 indicate PAHs source to be of petroleum origin while ratios larger than 0.1 indicate PAHs source to be of combustion origin and the ratio of (BaA / (BaA+Chr)) less than 0.2 implies petrogenic, from 0.2 to 0.35 indicates either petrogenic or pyrogenic origins, and larger than 0.35 implies pyrogenic sources [14].

(Flut/Pyr) = fluoranthene to pyrene.

(Phen/Ant) = Phenanthrene to Anthracene.

(Ant / (Ant + Phe)) = Anthracene to (Anthracene + Phenanthrene).

(BaP / (BaP + Chr)) = benzo (a) anthracene to (benzo (a) anthracene + chrysene).

LMW (low molecular weight) = Naphthalene + Acenaphthylene + Acenaphthene + Fluorene + Phenanthrene + Anthracene.

HMW (high molecular weight) = Fluoranthene + Pyrene + Benzo[a]anthracene + Chrysene + benzo(b)fluoranthene + Benzo[k]fluoranthene + benzo(a)pyrene + Dibenzo[a,h]anthracene + benzo(ghi)perylene + Indeno[1,2,3-cd] pyrene.

RESULTS AND DISCUSSION

From the results obtained levels (ng/g dry weight) of the sixteen PAHs distribution in *Lu. xanthopterus*, *Ct. idella*, *Cy. carpio*, *T. zillii*, *P. abu* and *Le. vorax* are presented in Tables 1. The total concentrations of PAHs in fishes vary from 0.02ng/g dry weights to 3.34ng/g dry weights (Figure 1). Shatt Al-Arab River is affected by hydrocarbons pollution by sewage, industrial wastes and the activities of refining in addition to natural seeps, and other human activities [15]. This may explain the difference that observed between the PAHs concentrations in the fish species.

The comparison revealed, elevated PAHs concentrations in Shatt AL-Arab River compared to those reported by [16] and [17]. It also indicated that Shatt Al-Arab River showed lower concentrations of PAHs compared to those found by [18,19,20,21,11,8,22].

Table 1: List of fish species included lipids content (%) in present study.

Fish species	No. Of fish	Total weight (g)	Total length (cm)	Fat%
<i>Lu. xanthopterus</i>	10	916	40	7.5
<i>Ct. idella</i>	18	1117	57	2.72
<i>Cy. carpio</i>	30	1019	35	2.6
<i>T. zillii</i>	60	75	15	2.5
<i>P. abu</i>	46	43	21	3.5
<i>Le. vorax</i>	30	750	35	2.5




Figure 1: Total concentrations of PAHs in some fishes of Shatt Al-Arab River.

The maximum concentration of PAHs in fish species was found in *L. vorax* and the minimum concentration of PAHs was found in *C. carpio*. (Table 2). The various components of PAHs in present studies Nap, Ace, Flu and Ant observed in *L. xanthopterus* (Figure 2). Residual levels of Acy, Ace, Flu, Phe, Ant, Flut, BaA, Chy and BbF were high in the tissues of *Ct.idella* (Figure 3). It is obvious from (Figure 4), that Nap, Ace, Phe, Pyr, BaA, Chyr and BbF are abundant

in *Cy. carpio*. The predominant components of PAHs in the *T. zillii* were ACY, Flu, Phe, Ant, Pyr, BaA and BPY (Figure 5). In the muscle of *P. abu*, the most abundant PAHs are Nap, Ace, Acy, Phe, Flue, BbF and BkF (Fig. 6). In the muscle of *Le. vorax* the most abundant PAHs are Ace, Acy, Flu, Flue, Pyr, BaA, Chy, BbF, BkF, BaP and DahA (Fig.7).

Table 2: Concentrations of Polycyclic Aromatic Hydrocarbons in the fish samples.

Name compound	<i>Lu. xanthopterus</i>	<i>Ct.idella</i>	<i>Cy. carpio</i>	<i>T. zillii</i>	<i>P. abu</i>	<i>Le. vorax</i>
Naphthalene	0.024		0.008		0.006	
Acenaphthylene		0.097		0.271	0.903	1.560
Acenaphthene	0.723	0.005	0.003		0.210	0.030
Fluorene	0.529	0.003		0.784		0.071
Phenanthrene		0.017	0.002	0.020	0.056	
Anthracene	1.824	0.082		0.396		
Fluoranthene		0.003			0.009	0.012
Pyrene			0.001	0.003	0.134	0.013
Benzo[a]anthracene		0.014	0.003	0.025		0.014
Chrysene		0.0004	0.001			0.005
Benzo[b]fluoranthene		0.002	0.002			0.002
Benzo[k]fluoranthene					0.523	0.034
Benzo[a]pyrene						0.014
Dibenzo[a,h]anthracene						1.588
Benzo[g,h,i]perylene				0.026		
Indeno[1,2,3-cd] pyrene						
Σ 16 PAHs	3.100	0.226	0.025	1.528	1.844	3.348

PAH isomeric ratios such as Flue/Pyr, Phe/Ant, Ant/(Ant+Phe), BaA/(BaA+Chry) and (LMW/HMW) low molecular weight to high molecular weight have been used to identify sources that contribute PAHs to the environment [23]. In the present study (Tables 3), the ratio of (Flu / Pyr) < 1 reflecting petrogenic origin, while the ratio of

(Phe/Ant) less than 1 indicates petrogenic. (Ant/(Ant+Phe)) ratio presumes that ratios larger than 0.1 indicate PAHs source to be of combustion origin and the ratio of (BaA/(BaA+Chr)) larger than 0.35 implies pyrogenic sources. ratio of (LMW /HMW) values less than 1 are attributed to pyrogenic sources[13].

Table 3. Values of PAHs ratios for the fishes of Shatt Al Arab River.

PAHs ratios	<i>Ct.idella</i>	<i>Cy. carpio</i>	<i>T. zillii</i>	<i>P. abu</i>	<i>Le. vorax</i>
fluoranthene/pyrene				0.070	0.956
Phen/Ant	0.217		0.051		
Ant/(Ant+Phen)	0.821		0.950		
BaA/(BaA+Chry)	0.972	0.669			0.717
LMW /HMW	12.058	1.473	26.889	1.802	1.000

In fact, the presence of fluoranthene and pyrene indicates the importance of pyrolytic inputs since these compounds are considered as products formed from the condensation of aromatic compounds of low molecular weight at high temperature [17]. The value of the Phen /Anth ratio in this

study is much lower than 10 which is generally considered indicative of a predominance of petrogenic sources [24]. The results of the present study indicated that fish were contaminated mainly with low-molecular weight (LMW) compounds, while the high-molecular weight (HMW) [25].




Figure 2: The percentage of the PAHs compound in *L. xanthopterus*




Figure 3: The percentage of the PAHs compound in *Ct. idella*




Figure 4: The percentage of the PAHs compound in *C. carpio*




Figure 5: The percentage of the PAHs compound in *T. Zillii*




Figure 6: The percentage of the PAHs compound in *P. abu*




Figure 7: The percentage of the PAHs compound in *Le. Vorax*

In this all the calculated values of the LMW/HMW ratio are >1 , and this indicates the main source of the PAHs in the investigated area coming from petrogenic source. LMW-PAHs are preferentially degraded during PAH transport and burial into sediments [26]. The ratios indicate that the main source of the PAHs, petrogenic input may also be a source of PAHs in these sites because many of the oil activities in the Shatt Al-Arab river that could contribute to PAHs [15].

The less carcinogenic polyaromatic hydrocarbon of lower molecular weight (LMW PAHs) was detected with clearly observed naphthalene and its substituent in all the studied fish species (Table 3). The more carcinogenic high molecular weight (HMW) PAHs (BaA, BbF, BkF, BaP, and InP) were low detected in the fish samples analyzed. This shows that the source of PAHs in the river was mainly from petrogenic sources. The ratio of low molecular weight PAHs (LMW-PAHs) to high molecular weight PAHs (HMW-PAHs) has been used to characterize the origin of PAHs in the environment [27]. The accidental discharge of hazardous materials such as petroleum and chemical solvents to the aquatic environment has become the focus of increasing regulatory and public concern because of the adverse impacts of such materials on human health and the environment [28].

Conclusion

Total concentrations of PAHs in the fishes were in the range of between 0.02 to 3.34 ng/g dry weight, which is considered by [29] as slightly polluted. Several ratios of PAHs species concentration were applied to identify sources of the anthropogenic pollutants, which indicate that the pyrolytic and petrogenic sources therefore should be preventing the various kinds of industrial and domestic contaminants to be discharged into the river.

REFERENCES

- Weiner, E.R. (2000). Application of Environmental chemistry. Lewis Publisher, London, New York.
- Tyagi, S.; Sharma, B.; Singh, P.; and Dobhal, R. (2013). Water Quality Assessment in Terms of Water Quality Index. *Amer. J. Wat. Res.*, 1(3):34-38.
- Al-Saad, H. T., Al-Hello A. A., Al-Kazaeh D. K., Al-Hello M. A., Hassan W. F., Mahdi S. (2015). Analysis of Water Quality Using Physico-Chemical Parameters in the Shatt Al-Arab Estuary, Iraq. *International Journal of Marine Science*, Vol.5, No.49: 1-9.
- UN-ESCWA and BGR (United Nations Economic and Social Commission for Western Asia; Bundesanstalt für Geowissenschaften und Rohstoffe), 2013, Inventory of Shared Water Resources in Western Beirut. CHAPTER 5 - SHATT AL ARAB, KARKHEH AND KARUN RIVERS.
- Wu, W.; Qin, N.; He, W.; He, Q.; Ouyang, H. and Xu, F. (2012). Levels, Distribution, and Health Risks of Polycyclic Aromatic Hydrocarbons in Four Freshwater Edible Fish Species from the Beijing Market. *The Scientific World Journal* Volume 2012, Article ID 156378, 12 pages.
- Azhari, A.; Dalimin, M.N.; and Wee, S. T. (2011). "Polycyclic Aromatic Hydrocarbons (PAHs) from Vehicle Emission in Vegetation of Highway Roadside in Johor, Malaysia". *Inter. J. Env. Sci. Devel.*, 2(6): 465-468.
- Al-Saad, H. T. (1995). "Distribution and sources of hydrocarbons in Shatt Al-Arab estuary and NW Arabian Gulf". Ph. D. thesis, Basrah University, 186 pp.
- Mahmoodi, M., Safahieh, A., Nikpour, Y. and Ghanemi K. (2012). Distribution and Sources of Polycyclic Aromatic Hydrocarbons in the Sediment of Bushehr Coastal Zone-Iran. *Iranica Journal of Energy & Environment* 3 (2): 173-179.
- Al-Khion, D. H.; Al-Ali, B. S.; Al-Nagar, G.; Al-Saad H. T.; Al-Anber, L. J.; Khwadem, A. A.; Saleh, S. M. and Zeidan, A.M. (2016). Polycyclic Aromatic Hydrocarbons in Some Fishes from the Iraqi Marine Waters. *Journal of Natural Sciences Research* Vol.6, No.10, p 2224-3186.
- Grimalt, J.O. & J. Oliver. 1993. Source input elucidation in aquatic systems by factor and principal component analysis of molecular marker data. *Anal. Chem. Acta.* 278:159-176. doi: 10.1016/0003-2670(93)80094-2
- Qiu, Y.W., G. Zhang, G.Q. Liu, L.L. Guo, X.D. Li and O. Wai. (2009). "Polycyclic aromatic hydrocarbons (PAHs) in the water column and sediment core of Deep Bay, South China". *Est. Coastal Shelf Sci.* 83:60-66. doi:10.1016/j.ecss.2009.03.018
- Rita, B.J. Peachey, 2003. Tributyltin and polycyclic aromatic hydrocarbon levels in Mobile Bay, Alabama: A review. *Marine Pollution Bulletin*, 46: 1365-1371.
- Zhu, L.; Chen, W.; Wang, J. and Shen, H. (2004). "Pollution survey of polycyclic aromatic hydrocarbons in surface water of Hazhou, China". *Chemosph.* 56: 1085 – 1095.
- Vrana, B., A. Pasch and P. Popp. (2001). "Polycyclic Aromatic hydrocarbon concentration and Patterns in sediments and surface water of Mansfeld region, Saxony". Anhalt, Germany. *J. Environ. Monit.* 3(6):602-609.
- Guo, W., M. He, Z. Yang, C. Lin, X. Quan and H. Wang (2007). "Distribution of polycyclic aromatic in water, suspended particulate matter and sediment from Daliao River watershed, China". *Chemosphere.* 68:93–104. Doi: 10.1016/j.chemosphere.2006.12.072
- Farid, W. A.; Al-Salman, A. N.; Ali, W. A.; Al-Saad, H. T.; Mahdi, S. and Al-Hello, A. A. (2016). Polycyclic Aromatic Hydrocarbons (PAHs) in the surface sediments of Shatt Al-Arab River, Basrah City, Southern Iraq. *Journal of Natural Sciences Research*. ISSN 2225-0921 (Online) Vol.6, No.8. 2224-3186 pp.
- Wang, C., Wang, W., He, S., Due, J. and Sun, Z. (2011). Sources and distribution of aliphatic and polycyclic aromatic hydrocarbons in Yellow River Delta Nature Reserve, China. *Applied Geochemistry*, 26: 1330-1336.
- Zrafi, I., Hizem, L., Chalghmi, H., Ghrabi, A., Rouabhi, M. and Saidane-Mosbahi, D. (2013). Aliphatic and aromatic biomarkers for petroleum

- hydrocarbon investigation in marine sediment. *Journal of Petroleum Science Research (JPSR)*, 2(4): 146-155.
18. De Luca, G., Furesi, A., Leardi, R., Micera, G., Panzanelli, A., Piu, P.C. and Sanna, G. (2004). Polycyclic aromatic hydrocarbons assessment in the sediments of the Porto Torres harbor (Northern Sardinia, Italy). *Marine Chemistry*, 86: 15–32.
 19. Trabelsi, S. and Driss, M. R. (2005). Polycyclic aromatic hydrocarbons in superficial coastal sediments from Bizerte Lagoon, Tunisia. *Marine Pollution Bulletin*, 50: 344-359.
 20. Bin, J., Hai-Long, Z., Guo-Qiang, H., Hui, D., Xin-Gang, L., Hong, Tu. S. and Rui, L. (2007). Characterization and distribution of polycyclic aromatic hydrocarbon in sediments of Haihe River, Tianjin, China. *Journal of Environmental Sciences*, 19: 306-311.
 21. Zrafi-Nouira, I., Khedir-Ghenim, Z., Zrafi, F., Bahri, R., Cheraief, I., Rouabhia, M., SaidaneMosbahi, D. (2008). Hydrocarbon pollution in the sediment from the Jarzouna-Bizerte coastal area of Tunisia (Mediterranean Sea). *Bulletin of Environmental Contamination and Toxicology*, 80: 566-572.
 22. Chen, Y.; Jia, R. and Yang, Sh.(2015). Distribution and Source of Polycyclic Aromatic Hydrocarbons (PAHs) in Water Dissolved Phase, Suspended Particulate Matter and Sediment from Weihe River in Northwest China *Int. J. Environ. Res. Public Health*, 12, 14148-14163.
 23. [Yu B, Xie X, Ma LQ, et al. 2013. Source, distribution and health risk assessment of polycyclic Aromatic Hydrocarbons in urban street dust from Tianjin. *Environ SciPollut Res*, doi: 10.1007/s11356-013-2190-2.
 24. Tolosa, I., Stephen de M., Mohammad R. S., Jean-Pierre.V., Jean. B and Chantal. C., 2003. Aliphatic and aromatic hydrocarbon in coastal Caspian Sea sediments. *Marine Pollution Bulletin* Vol. 48, Issues 1-2, January 2004, Pages 44-60. Trabelsi, S. and Driss, M. R. (2005). Polycyclic aromatic hydrocarbons in superficial coastal sediments from Bizerte Lagoon, Tunisia. *Marine Pollution Bulletin*, 50: 344-359.
 25. Malik, A., Ojha,P., Singh,K.P., 2008. Distribution of polycyclic aromatic hydrocarbons in edible fish from Gomti River, India. *Bull. Environ. Contam. Toxicol.* 80, 134–138.
 26. Berto, D., Cacciatore, F., Ausili, A., Sunseri, G., Luca, G., Bellucci, L. G., Frignani, M., Albertazzi, S. & Giani, M. (2009). Polycyclic Aromatic Hydrocarbons (PAHs) from Diffuse Sources in Coastal Sediments of a Not Industrialised Mediterranean Island. *Wat. Air Soil Pollut.* 200:199–209.
 27. Olaji, E. D. Nwogu, N. A. Yakubu, A. F. and Olaji, C. O. 2014. "Assessment of Total Hydrocarbon Concentration in Four Fish Species of Degele Community, Nigeria and Their Dietary Intake in the Populace ". *Advances in Research* 2(2): 109-118.
 28. Al-Shwafi, N.A.A. 2008. "Total Petroleum hydrocarbon carcinogens in commercial fish in the Red sea and Gulf of Aden-Yemen, JKAU: Mar.Sci. 15-28.
 29. Unlu, S., and Alpar, B., 2006 Distribution and Sources of Hydrocarbons in Surface Sediments of Gemlik Bay (Marmara Sea, Turkey), *Chemosphere* 64, 764-777.

Source of support: Nil; Conflict of interest: None declared