

**EXTRA CELLULAR ABILITY OF *BACILLUS SUBTILIS* AND
PSEUDOMONAS FLUORESCENS TO DEGRADE AFLATOXIN B1
PRODUCED BY *ASPERGILLUS FLAVUS***

Labeed A. Al-Saad^{*1}, Adnan I. Al-Badran², Sami A. Al-Jumayli³

¹University of Basrah, College of Agriculture, Dept. of Plant Protection.

²University of Basrah, College of sciences, Dept. of Biology.

³University of Karbala, College of Applied Medical Sciences.

Article Received on
29 May 2015,

Revised on 21 June 2015,
Accepted on 12 July 2015

***Correspondence for
Author**

Dr. Labeed A. Al-Saad

University of Basrah,
College of Agriculture,
Dept. of Plant Protection.

ABSTRACT

This study was designed to evaluate the extra cellular ability of *Bacillus subtilis* isolates (BSS1, BSS2, BSS3, BSS4 and BSW) and *Pseudomonas fluorescens* isolates (PFMst and PFDL) to degrade aflatoxin B1 (AFB1) that produced by *Aspergillus flavus* AFL14. The extra-cellular degradation ability of bacterial isolates to AFB1 was evaluated quantitatively using HPTLC technique. *B. subtilis* isolates BSS4, BSW and BSS3 were exceeded the rest of bacterial isolates significantly (100, 100 and 99.55) % respectively, followed by *P. fluorescens* isolates PFMst and PFDL (97.805 and 97.396) % respectively, with no significant differences among individuals in the

same group. The degradation residues administrated to rats to determine their effect on biosystems, the blood parameters showed a significant reduction in WBC, HB, RBC, and P.C.V while there was a significant increase in Urea, Glutamic pyrovate transaminase (GPT) and Glutamic oxaloacetic transaminase (GOT) in AF+DMSO treatment comparing with the negative and positive control which is may indicate that AFB1 caused immune-malfunction, hemolysis, kidney malfunction and liver stress. The biocontrol treatments showed a significant increase in WBC caused by BSS4+AF treatment, significant reduction in RBC caused by BSS4 and PFMst and significantly high Urea levels exceeded AF+DMSO treatment caused by PFMst and PFMst+AF treatments respectively. According to platelet count parameter, no significant differences were observes among all treatments and control treatments.

KEYWORDS: *Aspergillus flavus*, aflatoxin B1, extra-cellular, degradation, *Pseudomonas fluorescens*, *Bacillus subtilis*.

INTRODUCTION

Aspergillus flavus is a worldwide fungus, highly distributed in soil, water and air and possesses the ability to contaminate crops, processed foods and feedstuff (Vujanovic *et al.*, 2001). A broad spectrum of human diseases were reported to be caused by *A.flavus* which included human tissue infections which involved skin, oral mucosa, or subcutaneous tissues and hypersensitivity reactions in addition to producing Aflatoxins, especially AFB1 which is the most toxic and potent hepatocarcinogenic natural compound ever characterized (Hedayati *et al.*, 2007).

The control of fungal diseases was conventionally performed by chemical compounds which were mostly affected the environment and human and animal health in addition to economical costs as this type of control is usually expensive. However, since 1930 the scientists discovered interesting beneficial microbes with antifungal activities that could be employed in controlling plant fungal diseases (Cawoy *et al.*, 2011). Among those microorganisms, a nontoxigenic filamentous fungi like *A.flavus* and *A.parasiticus* strains (Rajani *et al.*, 2012), *Tricoderma harzianum* (Al-Othman *et al.*, 2013), yeasts like *Candida krusei*, *Saccharomyces cerevisiae* and *Pichia anomala* (Sari *et al.*, 2008; Rajani *et al.*, 2012) and bacteria like *Bacillus subtilis*, *Pseudomonas spp.*, *Ralstonia spp.* and *Lactobacilli spp.* (Mejía-Teniente *et al.*, 2011; Rajani *et al.*, 2012). *Bacillus subtilis* have some characteristics that enabled it to be highly recommended as a biological control agent as being recognized as non-pathogenic organism by the US Food and Drug Administration (USFDA), high tolerant to hard conditions like low nutrition and water supply, high temperature and unfavourable pH in addition to its ability to produce spores which is important benefit enabled ease of handling of this bacteria by converting to powder formulation (Cawoy *et al.*, 2011). *B.subtilis* have the ability to inhibit spore germination and *A.flavus* growth in addition to its extracellular activity to degrade more than 50-78.39% of aflatoxin B1 with maximum activity at 50°C and wide range of pH (5-8) (Farzaneha *et al.*, 2010; Thakaew and Niamsup, 2013; Watanakij, 2013). *P.fluorescens* is one of the important biocontrol agents found to inhibit *A.flavus* growth and AFB1 production and/or AFB1 content reduction either *in vitro* or *in vivo* (Luna-Romero *et al.*, 2000; Reddy *et al.* 2010; Singh *et al.*, 2011; Mohale, 2013). *P.fluorescens* possessed some characteristics helped to considering it as effective biocontrol agent that included the ability

to grow and rapid colonization of the rhizosphere and spermosphere in soil environment, aggressive competitor with high ability to adapt environmental stress, the ability to produce wide spectrum of extra-cellular enzymes (cellulase, chitinase, amylase, protease, and gelatinase), antibiotics, siderophore, growth promoter substances and volatiles, furthermore, each strain is able to act against the same pathogene in multiple patho-system (Weller, 2007; Couillerot *et al.*, 2009; Soesanto *et al.*, 2011). The Public Health Agency of Canada classified *P.fluorescens* as a risk group 1 which means that it can infect human in contact with contaminated medical devices, immunocompromised patients and people with outbreaks but it is unlikely infects general population or causes diseases to healthy research works (Canadian Dept. of Environment and Canadian Dept. of Health, 2013).

MATERIALS AND METHODS

Microorganisms: *A.flavus* AFL14 was isolated from soil using dilution method on MEA (Malt Extract Agar: Oxoid Limited) then preliminary identified morphologically according to Klich (2002) , Molecular identification was made to confirm isolates identification according to . Rodriguez *et al.* (2012) using Internal Transcript spacer ITS1-ITS2 primers:

(For: 5'TCCGTAGGTGAACCTGCGG3',

Rev:5' GCTGCGTTCTTCATCGATGC 3')

and ITS3-ITS4 region (For: 5'GCATCGATGAAGAACGCAGC3',

Rev: 5'TCCTCCGCTTATTGATATGC3') (Bellemain *et al.*, 2010)

Bacillus subtilis isolates BSS1, BSS2, BSS3, and BSS4 are obtained as a gift from Dr. Sami Al-Jumaily, Karbala Univ., College of Applied Medical Sciences; BSW is obtained as a gift from Dr. Wael Al-Waely, Basrah Univ., College of Agriculture, *Pseudomonas fluorescens* PFMst are obtained from Dr. Mohammed Amir; PFDL is obtained from Dr. Dehya Al-Waely, Basrah Univ., College of Agriculture, Plant Protection Dept.

All *Bacillus subtilis* isolates are confirmed morphologically and biochemically (Gram staining, shape, spore existence, oxidase test, catalase test, gelatin liquefaction, indol test, methyl red test, Voges-Proskauer test, citrate test, salt tolerance test and amylase test) whereas *Pseudomonas* isolates are confirmed by (Fluorescens under UV light, gram staining, shape, oxidase test, catalase test, gelatin liquefaction, indol test, methyl red test, Voges-Proskauer test) (Henayl, 2000; Barrow and Filtham, 2003; Forbes *et al.*, 2007).

The extra-cellular ability of bacterial isolates to degrade AFB1

Aflatoxin B1 was mixed with 10 ml of sterilized Nutrient broth medium at a concentration of 250 ng/ml. The medium was inoculated with 1 ml of (1×10^6 CFU) of 24h age bacterial broth culture of *B. subtilis* isolates BSS1, BSS2, BSS3, BSS4, BSW and *P. fluorescens* isolates PFMst and PFDL. Treated tubes were incubated at 35°C in shaking incubator for 48h (Zuo *et al.*, 2012).

High performance thin layer chromatography (HPTLC)

AFB1 was extracted according to Al-Jumaily (1996) and the qualitative TLC was obtained according to Goldblatt (1969). Quantitative estimation was obtained using 6 captures for each TLC plate from different positions using a digital camera (CASIO EXILIM EX-Z80, Japan) to achieve precision, the images were analyzed with Image J 1.46r open source software (National Institute of Health, USA: <http://imagej.nih.gov/ij>) to record fluorescens intensities of the AFB1 spots. A standard curve was made using linear plotting of AFB1 standard spots fluorescens (under UV light $\lambda=366$ NM) values versus AFB1 concentration (Hoeltz *et al.*, 2010; Hoeltz *et al.*, 2012). The experiment was designed in one way CRD and the statistical analysis was carried out by SPSS® Ver. 16.0 software.

Bioassay for AFB1 degradation residues

Two bacterial isolates, *B. subtilis* BSS4 and *P. fluorescens* PFMst were chosen according to the HPTLC results that showed a high ability to degrade AFB1 (approximately 100%) and to determine the effect of AFB1 degradation residues *in vivo*. The bacterial broth cultures, after incubation for 48h with 250 ng/ml of AFB1 were preceded with equal amounts of chloroform then filtrated with Whatman No. 1 filter paper using Buchner funnel. The extracts were concentrated to dryness in water bath at 40°C and the crystallized residues were re-dissolved with the physiological solution Dimethyl sulfoxid (DMSO) in addition to the standard positive treatment involved dissolving of AFB1 (250 ng/ml) in DMSO to be obtained in the oral injection of experimental animals. The bioassay is held using 28 albino rats, *Rattus rattus* of 8 weeks age. The animals were divided into 7 groups that were exposed to 7 different treatments. The treatments were (PFMst, PFMst+AFB1, BSS4, BSS4+AFB1, DMSO+AFB1, DMSO and non treated animals as a negative control), each treatment was obtained in 4 replicates. The treatments were performed by oral gavage of 4 doses equivalent to 250 ng/kg of body weight, with an interval of 2 days, after the last dose the animals left for one month then they were scarified through opining the abdominal alveolus by dissection after

anesthetization with Di-ethyl ether. Blood samples were collected by heart puncture technique and collected blood is divided into two groups, the first one placed in EDTA containing tubes for physiological tests that carried out using an Auto Hematology analyzer (ABX Micros ES60, France), whereas the second group is placed in EDTA free tubes for biochemical tests which is carried out using Reflotron®Plus devise (Roche Ltd., Germany) (Gagini *et al.*, 2010; El-Hady *et al.*, 2011; Al-Saadi, 2012). The experiment was designed in one way CRD and the statistical analysis is carried out by SPSS® ver. 16.0 software.

RESULTS AND DISCUSSION

The morphological identification test approved the identity of the isolate AFL14 as *A.flavus* which is confirmed by molecular identification results which revealed that 100% alignment between ITS1-ITS2 PCR product sequence with the standard isolate (CBS 100558; ID:gb|KJ175473.1) and 100% alignment between ITS3-ITS4 PCR product sequence with the standard isolate (M1204.653; ID:gb|KJ175474.1) in NCBI gene bank.

Bacterial isolate

The biochemical test results of *B. subtilis* and *P. fluorescens* isolates presented in tables (1 and 2) respectively were consistent with the same test results in Henayl, (2000), Barrow and Filtham (2003), Forbes *et al.* (2007) which confirmed the previous definition conducted in advance by the source donors.

Table (1): Biochemical tests of *B. subtilis*.

Isolate	Gram stain	Shape	Spores	Oxidase	Catalase	Gelatine	IMVC				NaCl 6.5%	Amelase
							Indol	MR	VP	Citrate		
BSS1	+	+	+	+	+	+	-	+	+	+	+	+
BSS2	+	+	+	+	+	+	-	+	+	+	+	+
BSS3	+	+	+	+	+	+	-	+	+	+	+	+
BSS4	+	+	+	+	+	+	-	+	+	+	+	+
BSW	+	+	+	+	+	+	-	+	+	+	+	+

Table (2): Biochemical tests of *P. fluorescens*.

Isolate	fluoresces under UV	Gram stain	shape	Oxidase	Catalase	Gelatine	IMVC			
							Indol	MR	VP	Citrate
PFMst	+	-	+	+	+	+	-	-	-	+
PFDL	+	-	+	+	+	+	-	-	-	+



Fig. 1: TLC plate of bacterial degradation of AFB1.

The quantitative analysis of TLC plates captures (Figure 1 and 2) revealed the high ability of all bacterial isolates to degrade AFB1 in liquid medium. The degradation percentage ranged from (95.56-100%) with a significant differences among isolates. BSS4, BSW and BSS3 exceeded the rest of isolates significantly with no significant differences among them followed by PFMst and PFDL which surpassed BSS1 whereas BSS2 possessed the lowest degradation ability. The extracellular degradation of AFB1 is mostly attributed to the extracellular enzymatic activity of the bacteria which is usually depended on the level of the genes that controlling the production of those enzymes (Reddy *et al.*, 2010; Afsharmanesh *et al.*, 2014) adding to that, there are many bacterial species (*B.subtilis*, *Escherichia.coli* and *Lactococcus lactis*) posses a high ability to detoxify AFB1 in liquid medium by binding AFB1 to bacterial cell wall due to the intimacy between them which is lead to detoxify aflatoxin B1 (Al-Jumaily, 2014).




Fig. 2: AFB1 degradation (%) by bacterial isolates. LSD=0.765 at $P \leq 0.05$

Bioassay for AFB1 degradation residues

Blood physiological measurements

The results of the several physiological parameters of blood were considered to determine the impact of AFB1 and its degradation side products on biological systems (rat was our model).

The parameters included:

White blood cells (WBCs)

The number of WBCs was influenced by the treatment derivatives that administered to the animals (Figure 3). AF+DMSO showed the lowest WBCs rate ($1.6 \times 10^9/L$) that was significantly differed from all other treatments, followed by (-) Control, PFMst, DMSO, PFMst+AF (7, 7.966, 9.6 and $10.35 \times 10^9/L$) respectively with no significant differences among them, in addition to significant increase in WBCs number in BSS4+AF ($14.375 \times 10^9/L$) treatment compared to AF+DMSO, (-) Control and PFMst (1.6, 7 and $7.966 \times 10^9/L$) respectively and similar increase in BSS4 ($12.6 \times 10^9/L$) compared to (AF+DMSO and (-) Control) with no significant differences between BSS4+AF and BSS4. Generally, the results revealed that WBCs levels were not affected by *P.fluorescens* in opposite to *B.subtilis* and its AFB1 degradation derivatives which caused WBCs levels to be doubled. This may be an indication that the immune system had not been affected in addition to that the bacterium itself may have antigens stimulated immune activity toward it.




Fig. 3: The effect of interaction of AFB1 x biocontrol agents on WBCs levels. LSD=4.204 at $P \leq 0.05$

The reduction in WBCs levels in AF+DMSO treatment could be returned to the deterioration of the immune system of the rat as AFB1 is well known as immunosuppressant (Mogilnaya *et al.*, 2010; Anokwuru *et al.*, 2011). The effect of AFB1 on the number of the WBCs found to be dose dependent due to stimulation or suppression of the immune system (Hinton *et al.*,

2003; Mogilnaya *et al.*, 2010; Anokwuru *et al.*, 2011). Our study corresponded with previous studies which reported a significant decrease in WBCs count due to AFB1 activity (El-Shewy and Ebrahim., 2004; Mogilnaya *et al.*, 2010; Omoya *et al.*, 2014).

Hemoglobin (HB)




Fig. 4: The effect of interaction of AFB1×biocontrol agents on HB levels. LSD=1.840 at $P \leq 0.05$

AF+DMSO treatment displayed a significant reduction level (12.533 g/L) compared to DMSO, BSS4+AF, (-) Control and BSS4 (14.433, 14.766, 14.966 and 15.1g/L), respectively with no significant differences with PFMst and PFMst+AF treatments (13.466 and 13.933 g/L) respectively (Figure 4). All other treatments had no significant differences among them. The reduction of HB could be caused as a result of oxidative damage due to the activity of AFB 1 which is reported to increase auto-oxidation rate of oxyhemoglobin in addition to haemolysis (Abdel-Wahhab *et al.*, 2014) or due to the binding of AFB1 with the cell membrane proteins which is affect the energy supply (ATP) to the red blood cells, as well as the transfer of sugar in to the cells which is eventually lead to shorten the cell life and death (Cloherty *et al.*, 2001) . Kubena *et al.*(1997) found that secondary metabolic products of *A.flavus* (aflatyoxins: B1, B2, G1and G2) have the ability to bind with blood proteins that are responsible for the biosynthesis of red blood cells leading to the lake in cell numbers which is affect the balance of the blood (Hemostasis).

Red blood cells (RBCs)

A significant reduction in RBCs number was observed in AF+DMSO treatment (Figure 5) compared to (BSS4+AF, DMSO and (-) Control) respectively, those treatments, on the other hand did not show a significant differences among them. BSS4 and PFMst also recorded a significant reduction in RBCs level compared to ((-) Control and DMSO) treatments, respectively while BSS4 reduction rate was significantly lower than (BSS4+AF, DMSO, (-)

Control) respectively. No significant differences observed among AF+DMSO, PFMst and BSS4 which were ordered respectively. RBCs reduction could be resulted in blood haemolysis that may be caused due to the potential of AFB1 (Abdel-Wahhab *et al.*, 2014).




Fig. 5: The effect of interaction of AFB1×biocontrol agents on RBCs levels. LSD=1.002 at $P \leq 0.05$

Hematocrit (HTC% or P.C.V.)

Figure (6) displayed that P.C.V. rate of AF+DMSO treatment was lower than all other treatments with significant differences compared to (DMSO, (-) Control, BSS4 and BSS4+AF) respectively and not significantly differed than (PFMst+AF and PFMst), respectively. No significant differences were observed among all other treatments. The percentage of red blood cells reduction may be related to the impact of AFB1 which causes chronic blood haemolysis that leads to cause folate deficiency that impaired the DNA synthesis and nuclear development causing anaemia which reduces P.C.V. level (Golder , 2007; Mandal *et al.*, 2013; Abdel-Wahhab *et al.*, 2014).




Fig. 6: The effect of interaction of AFB1×biocontrol agents on P.C.V. levels. LSD= 5.86 at $P \leq 0.05$

Platelet count (PLT)

The results of platelet count (Figure 7) showed no significant differences among positive and negative control comparing to the rest of experiment treatments included AF+DMSO treatment. The only significant difference was between BSS4+AF ($760.22 \times 10^9/L$) and PFMst ($496.67 \times 10^9/L$). These results were similar to Sharma *et al.* (2011) results on mice as they found no significant differences in platelet count in blood of mice when they exposed to 2ug/kg of body weight of AFB1 comparing to control treatment.




Fig. 7: The effect of interaction of AFB1×biocontrol agents on platelet count. LSD=252.03 at $P \leq 0.05$

Mean corpuscular volume (MCV)

The results in figure (8) showed that MCV parameter under AF+DMSO treatment (61.466 fL) was significantly exceeded all other treatments included (positive and negative) control treatments followed by PFMst and PFMst+AF (59.9667 and 59.2667 fL), respectively with no significant differences between them. BSS4+AF (52.667 fL) displayed no significant difference either with control or DMSO treatments. The increase in corpuscular volume known to be related to folic acid deficiency (Golder, 2007). This result accompanying with reduction in WBC, RBC, HB and P.C.V may led to believe that AFB1 caused a megaloplastic anemia to the treated rats (Golder, 2007).




Fig. 8: The effect of interaction of AFB1×biocontrol agents on MCV levels. LSD= 4.392 at P≤0.05

Blood biochemical measurements

Urea

A significant increases in urea levels were observed in all treatment compared to (-) control and DMSO treatments (26.133 and 31.933 mg/dl) respectively (Figure 9) giving an indication of renal malfunction as the rate of renal clearance becomes lower than blood urea production which in turn can be considered as nephrotoxicity indicator (Mandal *et al.*, 2013). AF+DMSO treatment (42.933 mg/dl) exceeded BSS4+AF treatment (33.666 mg/dl) significantly whereas it was retracted significantly against PFMst+AF and PFMst (46.066 and 45.133 mg/dl) respectively and that may be returned to the ability of *P.fluorescens* to produce toxic compound and/or that the degradation residues had a toxic potential on kidney. PFMst+AF, PFMst, BSS4 were surpassed BSS4+AF treatment significantly with no significant differences among them. Those results agreed with the past studies that reported a high urea levels after treating with AFB1 (El-Shewy and Ebrahim., 2004; Devendran and Balasubramanian, 2011).




Fig. 9: The effect of interaction of AFB1×biocontrol agents on Urea levels. LSD=7.408 at P≤0.05

Glutamic pyrovate transaminase (GPT) and Glutamic oxaloacetic transaminase (GOT)

The GPT results displayed in Figure (10) revealed the superiority of AF+DMSO (39.433 U/L) over all other treatments significantly followed by BSS4+AF (27.833 U/L) respectively while (Figure 11) demonstrated that GOT levels in AF+DMSO (402 U/L) treatment was exceeded all the other treatments. These results reflected the toxic effect of AFB1 to the liver tissue as it lyses the hepatic cells membrane that resulting in the releasing of GPT and GOT to the blood which leads to elevate their levels. Moreover, aflatoxins may affect other organs like kidney and endoplasmic reticulum that contained those enzymes which may lead to same results (Al-Saadi, 2012). The results of our study corresponded with the earlier studies that showed similar effect of AFB1 represented by the elevation of GPT and GOT levels (El-Shewy and Ebrahim., 2004; Mishra and Kishore, 2009; Al-Saadi, 2012).




Fig. 10: The effect of interaction of AFB1×biocontrol agents on GPT levels. LSD=6.302 at $P \leq 0.05$




Fig. 11: The effect of interaction of AFB1×biocontrol agents on GOT levels. LSD=39.129 at $P \leq 0.05$

REFERENCES

1. Abdel-Wahhab KG, Mannaa FA and Abdel-Wahhab MA. Panax ginseng C.A. Meyer Extract Protects Rat Erythrocyte from the Oxidative Damage Induced by the Synergistic Effects of Subchronic Treatment with Aflatoxin B1 and Fumonisin. *British Journal of Medicine & Medical Research.*, 2014; 4(9): 1883-1901.
2. Afsharmanesh H, Ahmadzadeh M, Javan-Nikkhah M and Behboudi K. 2014. Improvement in biocontrol activity of *Bacillus subtilis* UTB1 against *Aspergillus flavus* using gamma-irradiation. *Crop Protection.*, 2014; 60: 83-92.
3. Al-Jumaily SA. Integrated control of *Aspergillus flavus* and aflatoxin B1 contamination in peanut crop. PhD thesis. University of Baghdad; 1996.
4. Al-Jumaily SA. Mycotoxins. Iraq; Book house publisher, 2014. 423 p.
5. Al-Othman MR., Mahmoud MA and Abd El-Aziz1 ARM. Effectiveness of nontoxigenic *Aspergillus flavus* and *Trichoderma harzianum* as biocontrol agents on aflatoxin B1 producing by *Aspergillus flavus* isolated from Cashew. *Life Science Journal.*, 2013; 10(4): 1918-1922.
6. Al-Saadi EBK. Characterization of *Aspergillus spp.* Isolates that are Carrying the *aflR* Gene and the Capability of Limiting their Infection to Some Foods in Al-Najaf Al-Ashraf Markets. MSc. Thesis. University of Kufa., Iraq, 2012.
7. Anokwuru CP, Ezekiel CN, Alabi OA, Shallie PD and Haruna MT. Ameliorative Effects of Ethanolic Neem Extract on Dietary Aflatoxin Induced Hematological Damage and Hepatotoxicity in Mice. *Asian Journal of Medical Sciences.*, 2011; 3(6): 254-260.
8. Barrow GI and Feltham RKA.. *Cowan and Steel's Manual for the identification of medical bacteria.* Cambridge University Press, The Edinburgh Building, Cambridge CB2 2RU, UK. 2003. 331p.
9. Bellemain E, Carlsen T, Brochmann C, Coissac E, Taberlet P and Kausrud H. ITS as an environmental DNA barcode for fungi: an in silico approach reveals potential PCR biases. *BMC Microbiology.*, 2010; 10: 189.
10. Canadian Dept. of Environment and Canadian Dept. of Health. Screening Assessment for *Pseudomonas fluorescens* ATCC 13525. Canadian Environmental Protection Act, 1999. Archived Government Notices., 2013; 143(40).
11. Cawoy H, Bettiol W, Fickers P and Ongena M. *Bacillus*-Based Biological Control of Plant Diseases. In: Margarita Stoytcheva. *Pesticides in the Modern World - Pesticides Use and Management.* InTech. Croatia. 2011, pp. 273-302.

12. Cloherty EK, Livine KB and Carruthers A. The red blood cell glucose transporter presents multiple nucleotide sensitive sugar exit sites. *Biochemistry.*, 2001; 40: 15549-15561.
13. Couillerot OC. Prigent-Combaret J. Caballero-Mellado and Moe nne-Loccoz Y. *Pseudomonas fluorescens* and closely-related fluorescent pseudomonads as biocontrol agents of soil-borne phytopathogens. *Letters in Applied Microbiology.*, 2009; 48: 505–512.
14. Devendran G. and Balasubramanian U. Biochemical and histopathological analysis of aflatoxin induced toxicity in liver and kidney of rat. *Asian Journal of Plant Science and Research.*, 2011; 1(4): 61-69.
15. El-Hady KA, Abdel-Hamid AK, Ibrahim A and Elzeftawy AE. Genotoxic And Cytotoxic Biomarkers Of Aflatoxin B1 And The Use Of Curcumin As Antidotal Therapy. *Zag. Vet. J.* (ISSN. 1110-1458). 2011; 39 (1): 86-102.
16. El-Shewy EA and Ebrahim MF. Ameliorative Effect Of Vitamin E Against The Toxicity Of Aflatoxin B1 On Rats With Special Reference To Its Effect On Male Fertility. 1st Ann. Confr. , FVM., Moshtohor., 2004; pp. 198-213.
17. Farzaneha M, Shi Z, Ghassempourc A, Ahmadzadeha M, Mirabolfathyd M, Javan-Nikkhaha M and Sharifi-Tehrani A. Evaluation of the potential of Iranian *Bacillus subtilis* strains to reduce aflatoxin B1 in pistachio by HPTLC and LC/MS/MS. *International Mycotoxin Conference - MycoRed*. Penang, Malaysia. 2010, p. 115.
18. Forbes BA, Sahm DF and Weissfeld AS. *Bailey and Acott's Diagnostic Microbiology*. Mosby, Inc., an affiliate of Elsevier Inc.China. 2007, 1031 pp.
19. Gagini TB, Silva RE, Castro IS, Soares BA, Lima MEF, Brito MF, Mazur C, Direito GM and Danelli MM. Oral Administration Of Piperine For The Control Of Aflatoxin Intoxication In Rats. *Brazilian Journal of Microbiology.*, 2010; 41: 345-348.
20. Goldblatt LA. *Aflatoxin: Scientific Background Control and Implication*. Food Science and Technology , A series of Monographs . Academic Press. New York and London. 1969. 472 p.
21. Golder B. 2007. The Anemias. in: Kligman RM, Jenson HB, Behrman RE and Stanton BF. *Nelson textbook of pediatric*.Saunders Elsevier. USA., 2007, 2005 p.
22. Hedayati MT., Pasqualotto AC, Warn PA, Bowyer P and Denning DW. *Aspergillus flavus*: human pathogen, allergen and mycotoxin producer. *Microbiology.*, 2007; 153: 1677-1692.

23. Henayl WR. Bergey's Manual of Determinative Bacteriology. Lippincott Williams & Wilkins. USA. 2000. 599 p.
24. Hinton DM., Myers MJ, Raybourne RA, Francke-Carroll S, Sotomayor RE, Shaddock J, Warbritton A and Chou MW. Immunotoxicity of Aflatoxin B1 in Rats: Effects on Lymphocytes and the Inflammatory Response in a Chronic Intermittent Dosing Study. *Toxicological Sciences.*, 2003; 73: 362–377.
25. Hoeltz M, Welke JE and Noll IB. Photometric Procedure For Quantitative Analysis Of Aflatoxin B1 In Peanuts By Thin-Layer Chromatography Using Charge Coupled Device Detector. *Quim. Nova.*, 2010; 33(1): 43-47.
26. Hoeltz M, Einloft TC, Oldoni VP, Dottori HA and Noll IB. The Occurrence of Aflatoxin B1 Contamination in Peanuts and Peanut Products Marketed in Southern Brazil. *Braz. Arch. Biol. Technol.*, 2012; 55(2): 313-317.
27. Klich MA. Identification of common *Aspergillus* species. Centraalbureau voor Schimmelcultuur, UTRECHT, The Netherlands. 2002. 116 p.
28. Kubena LF, Harvey RB, Corrier DE and Huff WE. Effect of feeding contaminated with DON and Vomitoxin in female of white leghorn chickens from day old through egg production. *Poult. Sci.*, 1997; 66: 1612-1618.
29. Luna-Romero II, Carvajal-Moreno M, Flores-Martínez A and Ferrera-Cerrato R. Possibility of Biological Control of *Aspergillus flavus* with *Pseudomonas fluorescens* on Maize Ear. *Revista Mexicana de FITOPATOLOGIA.*, 2000; 52.(18): 50-54.
30. Mandal A, Das K, Roy S, Mondal KC, Nandi DK. In vivo assessment of bacteriotherapy on acetaminophen-induced uremic rats. *JNEPHROL.*, 2013; 26(1): 228-236.
31. Mejía-Teniente L, Chapa-Oliver AM, Vazquez-Cruz MA, Torres-Pacheco I and Guevara-González RG. in: Torres-Pacheco I. Aflatoxins - Detection, Measurement and Control. In Tech. Croatia. 2011, pp. 317-354.
32. Mishra KS and Kishore K. Protective Effects of *Cichoricum intybus* Linn (Kasni) Against Dietary Aflatoxicosis in White Albino Rats. *American-Eurasian Journal of Toxicological Sciences.*, 2009; 1(1): 26-31.
33. Mogilnaya OA, Puzyr AP, Baron AV and Bondar VS. Hematological Parameters and the State of Liver Cells of Rats After Oral Administration of Aflatoxin B1 Alone and Together with Nanodiamonds. *Nanoscale Res. Lett.*, 2010; 5: 908–912.
34. Mohale SA. Impact of Ecophysiological Factors on Biocontrol of Aflatoxin Contamination of Maize by Atoxigenic *Aspergillus flavus* Strains. PhD.Thesis. Cranfield University. UK. 2013.

35. Omoya FO, Boboye B and Akinyosoye FA. Physiological Changes in Wister Rats Induced with *Bacillus* Species Used as Bio Control Agent. *Annual Research & Review in Biology.*, 2014; 4(11): 1798-1812.
36. Rajani P, Sridevi V and Lakshmi MVVC. A Review on Biological Control of Aflatoxin Crop Contamination. *International Journal of Chemical, Environmental and Pharmaceutical Research.*, 2012; 3(1): 83-86.
37. Reddy KRN, Raghavender CR, Reddy BN and Salleh B. Biological control of *Aspergillus flavus* growth and subsequent aflatoxin B1 production in sorghum grains. *African Journal of Biotechnology.*, 2010; 9(27): 4247- 4250.
38. Rodríguez A, Rodríguez M, Luque MI, Justesen AF, Córdoba JJ. A comparative study of DNA extraction methods to be used in real-time PCR based quantification of ochratoxin A-producing molds in food products. *Food Control.* 25: 666-672. aflatoxin B1 production in sorghum grains. *African Journal of Biotechnology.*, 2012; 9(27): 4247-4250.
39. Sari L., Skrinjar MM, Saka MB, Plavši DV and Abarkapa IS. Effect of the baker's yeast on the production of aflatoxin B1 by *Aspergillus flavus*. *Food Processing, Quality and Safety.*, 2008; 4: 165-168.
40. Sharma V, Sharma C and Sharma S. Influence of *Curcuma longa* and Curcumin on Blood Profile in Mice Subjected to Aflatoxin B1. *International Journal of Pharmaceutical Science and Research.*, 2011; 2(7): 1740-1745.
41. Singh SK, Sheeba RD, Gupta RS, Verma SK, Siddiqui MA, Mathur A, Agarwal PK Assessment of the Role of *Pseudomonas fluorescens* as Biocontrol Agent against Fungal Plant Pathogens. *Current Botany.*, 2011; 2(3): 43-46.
42. Soesanto L, Mugiastuti E, and Rahayuniati RF. Biochemical Characteristic Of *Pseudomonas fluorescens* P60. *Journal of Biotechnology and Biodiversity.*, 2011; 2: 19-26.
43. Thakaew R and Niamsup H. Inhibitory Activity of *Bacillus subtilis* BCC 6327 Metabolites against Growth of Aflatoxigenic Fungi Isolated from Bird Chili Powder. *International Journal of Bioscience, Biochemistry and Bioinformatics.*, 2013; 3(1): 27-32.
44. Vujanovic V, Smoragiewicz W and Krzysztyniak K. Airborne fungal ecological niche determination as one of the possibilities for indirect mycotoxin risk assessment in indoor air. *Environ Toxicol.*, 2001; 16: 1–8.

45. Watanakij N, Petchkongkaew A and Visessanguan W. Localization and Characterization of Aflatoxin B1-degrading Activity of *Bacillus subtilis* IRU-1A. The 15th Food Innovation Asia Conference. BITEC Bangna, Bangkok, Thailand. 2013.
46. Weller DM. *Pseudomonas* biocontrol agents of soilborne pathogens: Looking back over 30 years. *Phytopathology.*, 2007; 97: 250-256.
47. Zuo R, Chang J, Yin Q, Wang P, Cheng W, Wang X, Liu J and Zheng Q. Inhibiting *Aspergillus flavus* growth and degrading aflatoxin B1 by combined beneficial microbes. *African Journal of Biotechnology.*, 2012; 11(65): 12903-12909.