



Available online at: <http://www.basra-science-journal.org>

ISSN –1817 –2695



Received 22-4-2015, Accepted 27-10-2015

## **On Truncated of classical Beta Operators**

Ali J.Mohammad<sup>1</sup>, Hanadi A. AbdulSatter<sup>2</sup> and Haneen J. Sadiq<sup>3</sup>

*University of Basra, College of Education for Pure Science, Dept. of Maths., Basra, Iraq<sup>1,2</sup>.*  
*University of Basra, College of Science, Dept. of Maths., Basra, Iraq<sup>3</sup>.*

<sup>1</sup>[alijasmoh@yahoo.com](mailto:alijasmoh@yahoo.com), <sup>2</sup>[habdulla07@gmail.com](mailto:habdulla07@gmail.com), <sup>3</sup>[haneen.mahdi@yahoo.com](mailto:haneen.mahdi@yahoo.com)

### **Abstract**

In this paper, we study the truncated of classical Beta operators  $\beta_{n,N}(x)$  and the error occurs by the approximation. We estimate the truncated error in terms of modulus of continuity and the weighted norm of the function being approximated.

**Keywords:** MKZ Truncated operators, approximated order, Beta operators, modulus of continuity.

**Introduction**

A study of truncated of sequence of linear positive operators is a branch of approximation theory. In [4],[9],[12],[11] and [5] there is a review of truncated of some especially Szász –Mirakyan operators. Here, we study the truncated of the sequences of linear positive operators (Beta operators  $\beta_{n,N}(x)$ ). Then, we find the

error occurs by this approximation of truncated of Beta operator in terms of modulus of continuity of a function being approximated. For more properties of Beta operators you can read the articles [1],[2],[3],[6] and [7].

In the beginning, we give some preliminaries as follow:

$$\beta_n(f; x) = \frac{1}{n} \sum_{k=0}^{\infty} b_{n,k}(x) f\left(\frac{k}{n}\right), \tag{1}$$

where

$$b_{n,k}(x) := n \binom{n+k}{k} x^k (1+x)^{-n-k-1} \tag{2}$$

$n \in \mathbb{N}$  and  $x \in \mathbb{R}_0 = [0, \infty)$ .

For the functions  $f$  belonging to the weighted space  $C_p$ ,

$$C_p = \{f \in C[0, \infty) : f(x) = O(1+x^p) \text{ for some } p \in \mathbb{N}\},$$

$$\|f\|_p \equiv \|f(\cdot)\|_p = \sup_{x \in \mathbb{R}_0} u_p(x) |f(x)|,$$

$$u_p(x) \beta_n(|f(t) - f(x)|; x) \leq K_1(p) \omega_2\left(f; C_p; \sqrt{\frac{x(1+x)}{n} + \frac{x(2x+1)}{n^2}}\right), \tag{3}$$

for  $f \in C_p, n \in \mathbb{N}, x \in \mathbb{R}_0$  where  $\omega_2(f; C_p)$  is the modulus of continuity of  $f$ , we can get the properties of  $\beta_{n,N}(f)$  as follows:

$$\beta_{n,N}(f; x) = \sum_{k=0}^N b_{n,k}(x) f\left(\frac{k}{n}\right).$$

We can evaluate that:

$$\sum_{k=0}^{\infty} b_{n,k}(x) = 1, \text{ for } x \in [0,1), n \in \mathbb{N}, \tag{4}$$

$$\beta_n(f(t); 0) = f(0) = \beta_{n,N}(f(t); 0)$$

and

$$\beta_{n,N}(f(t); x) - f(x) = \beta_{n,N}(f(t) - f(x); x) - f(x) \sum_{k=N+1}^{\infty} b_{n,k}(x) \tag{5}$$

for every  $f \in C(I), x \in (0,1)$  and  $n, N \in \mathbb{N}$ .

In some papers ([4],[9],[11]and[12]) the authors investigated the truncated Szász-Mirakyan operators defined as:

$$S_{n,N}(f; x) = e^{-nx} \sum_{k=0}^N \frac{(nx)^k}{k!} f\left(\frac{k}{n}\right), \quad x \in \mathbb{R}_0, n \in \mathbb{N},$$

where  $f: \mathbb{R}_0 \rightarrow \mathbb{R}$ , and  $N = N(n, x)$  are positive integers depending on  $n$  and  $x$ . The results presented in some papers [4],[9]and [12] show that if  $x > 0$  is a fixed point and  $N = N(n, x)$  are integers such that  $N > nx$  for  $n \in \mathbb{N}$  and  $\lim_{n \rightarrow \infty} \frac{N-nx}{\sqrt{n}} = \infty$ , then

$$\lim_{n \rightarrow \infty} S_{n,N}(f; x) = f(x) \text{ for every } f \in C_p, p \in \mathbb{N}_0.$$

The aim of this note is to derive similar results for the truncated MKZ operators.

L. Rempulska and M. Skorupka in [10] are considered a strong approximation of  $f \in C_p$  for some linear positive operators.

The well-known Meyer-König and Zeller operators that we called MKZ is given by

$$M_n(f; x) = \begin{cases} \sum_{k=0}^{\infty} p_{n,k}(x) f\left(\frac{k}{n+k}\right) & \text{if } 0 \leq x < 1 \\ f(1) & \text{if } x = 1 \end{cases} \quad (6)$$

$$p_{n,k}(x) = \binom{n+k}{k} x^k (1-x)^{n+1} \quad (7)$$

Where  $f \in C(I), x \in I = [0,1]$  and  $n \in \mathbb{N} = \{1,2, \dots\}, C(I)$  is the space of continuous functions on  $I$  with the following norm

$$\|f\| = \sup\{|f(x)|: x \in I\}$$

$$\text{and } \|M_n(f) - f\| \leq \frac{31}{27} \omega\left(f, \frac{1}{\sqrt{n}}\right), n \in \mathbb{N}. \quad (8)$$

In [10] Rempulska were examined the strong approximation of by MKZ operators and he proved the inequality

$$M_n(|f(t) - f(x)|; x) \leq \frac{3}{2} \omega\left(f; \frac{1}{\sqrt{n}}\right) \quad (9)$$

for  $x \in [0,1], n \in \mathbb{N}$ .

In this paper, we show that an analog of MKZ operators connected with Beta operators.

**Theorem (1).** [5]

Let  $x \in (0,1)$  and  $n \in \mathbb{N}$  be fixed, and let  $N = N(n, x)$  be an integer such that

$N > (n + 1) \frac{x}{1-x}$ . Then

$$|M_{n,N}(f(t); x) - f(x)| \leq \frac{3}{2} \omega\left(f; \frac{1}{\sqrt{n}}\right) + |f(x)| \frac{1}{\sqrt{2\pi nx}} \frac{N}{(1-x)N-nx+1-x} \quad (10)$$

holds for every  $f \in C(I)$ .

**Corollary(1).** [5]

Let  $x_0 \in (0,1)$  and let  $N = N(n, x_0)$  be integers such that

- (i)  $N > \frac{(n+1)x_0}{(1-x_0)}$  for  $n \in \mathbb{N}$ ,
- (ii)  $(N/n)_{n=1}^{\infty}$  is a bounded sequence,
- (iii)  $\lim_{n \rightarrow \infty} \frac{N - \frac{(n+1)x_0}{1-x_0}}{\sqrt{n}} = \infty$

Then

$$\lim_{n \rightarrow \infty} M_{n,N}(f; x_0) = f(x_0) \text{ holds for every } f \in C(I). \tag{11}$$

**Corollary(2).** [5]

Let  $x_0 \in (0,1)$  and let  $N = \left[ \frac{n+x_0}{1-x_0} \right]$  for  $n \in \mathbb{N}$ . ( $[y]$  denotes the integral part of  $y \in \mathbb{R}$ ). Then the conditions (i)-(iii) are satisfied, and consequently the convergence (8) holds for every  $f \in C(I)$ . Moreover, we have

$$|M_{n,N}(f; x_0) - f(x_0)| \leq \frac{3}{2} \omega \left( f; \frac{1}{\sqrt{n}} \right) + K_2(x_0) |f(x_0)| \frac{1}{\sqrt{n}}$$

for  $f \in C(I)$  and  $n \in \mathbb{N}$ , where  $K_2(x_0) = \text{const.} > 0$ .

**Corollary(3).** [5]

If  $x_0 \in (0,1)$  and  $N = \left[ (n+1) \frac{x_0}{1-x_0} + \sqrt{n\alpha_n} \right]$  for  $n \in \mathbb{N}$ , where  $(\alpha_n)$  is a non- bounded sequence of positive numbers such that  $\left( \frac{\alpha_n}{\sqrt{n}} \right)_1^{\infty}$  is bounded , then (i)-(iii) and

(11) are satisfied , and the inequality

$$|M_{n,N}(f; x_0) - f(x_0)| \leq \frac{3}{2} \omega \left( f; \frac{1}{\sqrt{n}} \right) + K_3(x_0) |f(x_0)| \frac{1}{\alpha_n}$$

holds for every  $x f \in C(I)$  and  $n \in \mathbb{N}$ , where  $K_3(x_0) = \text{const.} > 0$ .

Here, we want to prove the above results in the space  $C_p$  with weighted norm:

**Theorem (2).**

Suppose that  $p \in \mathbb{N}_0$ ,  $n \in \mathbb{N}$  and  $x \in (0, \infty)$ . For  $N = N(n, x)$  be an integer such that  $N > nx$  , then the inequality

$$u_p(x) |\beta_{n,N}(f(t, x) - f(x); x)| \leq K_1(p) \omega_2 \left( f; C_p; \sqrt{\frac{x(1+x)}{n} + \frac{x(2x+1)}{n^2}} \right) + \|f\|_p \frac{\sqrt{n(1+x)+1} (x+1)N}{\sqrt{2\pi(n^2x-nx)} N+nx}, \tag{12}$$

holds for every  $f \in C_p$ , where  $K_1(p) > 0$  is constant and is given in (3).

**Proof:**

For given,  $f \in C(I), x \in (0,1)$  and  $n, N \in \mathbb{N}$  we get by (1),(3),(4) and (5):

$$\beta_{n,N}(f(t); x) - f(x) = \beta_{n,N}(f(t) - f(x); x) - f(x) \tag{13}$$

$$u_p(x) |\beta_{n,N}(f(t); x) - f(x)|$$

$$\begin{aligned} &\leq u_p(x) \beta_{n,N}(|f(t) - f(x)|; x) + \left| u_p(x) f(t) \sum_{k=N+1}^{\infty} b_{n,k}(x) \right| \\ &\leq K_1(p) \omega \left( f; C_p; \sqrt{\frac{x(x+1)}{n} + \frac{x(2x+1)}{n^2}} \right) + u_p(x) |f(x)| \sum_{k=1}^{\infty} b_{n,N+k}(x) \\ &\leq K_1(p) \omega \left( f; C_p; \sqrt{\frac{x(x+1)}{n} + \frac{x(2x+1)}{n^2}} \right) \\ &\quad + u_p(x) |f(x)| n(1+x)^{-n-1} \sum_{k=1}^{\infty} \binom{n+N+k}{N+k} x^{N+k} (1+x)^{-k} \\ &\leq K_1(p) \omega \left( f; C_p; \sqrt{\frac{x(x+1)}{n} + \frac{x(2x+1)}{n^2}} \right) \\ &\quad + u_p(x) |f(x)| b_{n,N}(x) \sum_{k=1}^{\infty} \frac{(n+N+1)(n+N+2) \dots (n+N+k)}{(N+1)(N+2) \dots (N+k)} \left(\frac{x}{1+x}\right)^k \\ &\leq K_1(p) \omega \left( f; C_p; \sqrt{\frac{x(x+1)}{n} + \frac{x(2x+1)}{n^2}} \right) \\ &\quad + u_p(x) |f(x)| b_{n,N}(x) \sum_{k=1}^{\infty} \left(1 + \frac{n}{N+1}\right) \left(1 + \frac{n}{N+2}\right) \dots \left(1 + \frac{n}{N+k}\right) \left(\frac{x}{1+x}\right)^k \\ &\leq K_1(p) \omega \left( f; C_p; \sqrt{\frac{x(x+1)}{n} + \frac{x(2x+1)}{n^2}} \right) \\ &\quad + u_p(x) |f(x)| b_{n,N}(x) \sum_{k=1}^{\infty} \left[ \left(1 + \frac{n}{N+1}\right) \frac{x}{1+x} \right]^k. \tag{14} \end{aligned}$$

By the assumptions:  $x \in (0,1)$  and  $N > nx$  we can write

$$\therefore \sum_{k=N+1}^{\infty} b_{n,k}(x) \leq b_{n,N}(x) \sum_{k=1}^{\infty} \left( \left(1 + \frac{n}{N+1}\right) \frac{x}{1+x} \right)^k$$

$$\begin{aligned} &< b_{n,N}(x) \sum_{k=1}^{\infty} \left( \left(1 + \frac{n}{N}\right) \frac{x}{1+x} \right)^k \\ &= b_{n,N}(x) \frac{(N+n)x}{N+nx}. \end{aligned}$$

If  $N > nx$  and  $x > 0$ , we have

$$\begin{aligned} b_{n,N}(x) &< b_{n,N} \left( \frac{N}{n} \right) = n \binom{n+N}{N} \left( \frac{N}{n} \right)^N \left( 1 + \frac{N}{n} \right)^{-n-N-1} \\ &= \frac{n(n+N)!}{N!(n-1)!} \frac{N^N e^{-N}}{N!} \left( \frac{(n+N)^{-n-N-1} e^{-n-N-1}}{(n+N)!} \right)^{-1} \left( \frac{n^n e^{-n-1}}{n!} \right). \end{aligned}$$

By using the Stirling formula, consequently

$n! \cong \sqrt{2\pi n} \left( \frac{n}{e} \right)^n$ , we have

$$\begin{aligned} b_{n,N}(x) &< \frac{n+1}{N+n+1} \left( \frac{1}{\sqrt{2\pi N}} \right)^{-1} \left( \frac{1}{\sqrt{2\pi(n-1)}} \right)^{-1} \frac{1}{\sqrt{2\pi(n+N+1)}} \\ &= \frac{\sqrt{2\pi(n+N+1)}}{\sqrt{4\pi N(n-1)}} = \frac{\sqrt{n+N+1}}{\sqrt{2\pi N(n-1)}} \end{aligned}$$

Therefore,

$$u_p(x) |f(x)| \sum_{k=N+1}^{\infty} b_{n,k}(x) \leq \|f\|_p \frac{\sqrt{n+N+1}}{\sqrt{2\pi N(n-1)}} \frac{(N+n)x}{N+nx}. \quad (15)$$

Putting  $N = nx$  then we get  $x = \frac{N}{n}$ ,  $x > 0$

$$\begin{aligned} u_p(x) |f(x)| \sum_{k=N+1}^{\infty} b_{n,k}(x) &\leq \|f\|_p \frac{\sqrt{n+nx+1}}{\sqrt{2\pi nx(n-1)}} \frac{(xn+n)N}{(N+nx)n} \\ &\leq \|f\|_p \frac{\sqrt{n(1+x)+1}}{\sqrt{2\pi(n^2x-nx)}} \frac{(xn+n)N}{(N+nx)n} \\ &\leq \|f\|_p \frac{\sqrt{n(1+x)+1}}{\sqrt{2\pi(n^2x-nx)}} \frac{(x+1)N}{(N+nx)}. \end{aligned}$$

from ( 13 ), ( 14 ), ( 15 ) we have results ( 12 ) ■

**Corollary(4).**

Let  $x_0 \in (0, \infty)$  and let  $N = (n, x_0)$  be positive integers such that for every  $n \in \mathbb{N}$ :

- (i)  $N > nx_0$ ,
- (ii)  $(N/n)_{n=1}^{\infty}$  is a bounded sequence,

$$(iii) \quad \lim_{n \rightarrow \infty} \frac{N-nx}{\sqrt{n}} = \infty.$$

Then the convergence

$$\lim_{n \rightarrow \infty} \beta_{n,N}(f; x_0) = f(x_0) \tag{16}$$

holds for every  $f \in C_p, p \in \mathbb{N}_0$ .

**Corollary(5)**

For a fixed  $x_0 > 0$  let  $N = [n(x_0 + 1)]$  for  $n \in \mathbb{N}$ . Then the conditions (i), (ii) and (iii) are satisfied, and hence (16) holds for  $f \in C_p$ . Also, we have

$$u_p(x_0) |\beta_{n,N}(f; x_0) - f(x_0)| \leq K_4(\omega(f; C_\rho; \sqrt{\frac{x_0^2 + x_0}{n} + \frac{x_0(2x_0 + 1)}{n^2}} + \|f\|_\rho \frac{1}{\sqrt{n}}))$$

for every  $f \in C_p$  and  $n \in \mathbb{N}$ , where  $K_4 = K_4(p, x_0)$  is a positive constant depending on  $\rho$  and  $x_0$ .

**Corollary(6)**

Let  $x_0 \in (0, \infty)$  and let  $N = [nx_0 + \sqrt{n}\alpha_n]$  for  $n \in \mathbb{N}$ , where  $(\alpha_n)$  is a non-bounded sequence of numbers  $\alpha_n \geq 1$  such that  $(\frac{\alpha_n}{\sqrt{n}})_{n=1}^\infty$  is bounded. Then (i), (ii) and (iii) are satisfied and hence (12) holds for every  $f \in C_p$ . Moreover, we have

$$\begin{aligned} u_p(x) |\beta_{n,N}(f; x_0) - f(x_0)| \\ \leq K_5(\rho, x_0)(\omega(f; C_\rho; \sqrt{\frac{x_0^2 + x_0}{n} + \frac{x_0(2x_0 + 1)}{n^2}}, \frac{1}{\alpha_n}) + \|f\|_\rho) \end{aligned}$$

for  $n \in \mathbb{N}$ , where  $K_5 = K_5(p, x_0) = \text{constant} > 0$ .

There are similar conditions of convergence of partial sums of truncated MKZ operators and Beta operator from theorem (1) and theorem (2). In our opinion, this is an expected result because both sequences are convergent to the function as  $n \rightarrow \infty$ .

## References

- [1] Vijay Gupta and A. Lupas, *On direct results for mixed Beta- Szász type operator*, Math. Vol. **13**, No.2 (2005) 83-94.
- [2] Vijay Gupta and Ogun Dođrn, *On direct results for Certain family of integral type-operators*, Journal of Inequalities in Pure and Applied Math. Vol. **6**, Issue 1, Article 21, 2005.
- [3] Vijay Gupta, Muhammad A., Man S., and M.K. Gupta, *On simultaneous approximation for certain Baskakov Durrmeyer type operators*, Journals of Inequalities in Pure and Applied Math. Volume **7**, Issue 4, Article 125 (2006).
- [4] H. G. Lehnhoff, *On a modified of Szász Mirakyan operator*, J.Approx. Theory,42 (1984), 278-282.
- [5] Lucyna Rempulska, *On truncated MKZ and Baskakov operators*, Journal of Math. Vol.6,(2012) no.8,377-384.
- [6] Prerna Maheshwari, *On some direct estimates for Szász-Beta operators in  $L_p$ -norm*, International Journal of Pure and Applied Math. S. Vol. **3** No.1 (2006) 61-72.
- [7] Prerna Maheshwari, *On saturation theorem for combination of modified Beta operators in  $L_p$ -spaces*, Applied Math. E-notes, **7** (2007) 32-37.
- [8] W. Meyer-König and K. Zeller, *Bernsteiniche Potenzreihen*, Studia Math. **19**(1960), 89-94.
- [9] E.Omey, *Note on operators of Szász -Mirakyan type*, J. Approx. Theory, **47**(1986),246-254.
- [10] L. Rempulska and M. Skorupka, *On strong approximation by modified Meyer- König and Zeller operators*, Tamkang J.Math., **37**(2), (2006),123-130.
- [11] Lucyna Rempulska, *On convergence of partial sums of operators of Szász -Mirakyan type*, International journal of pure and applied mathematics,**77**(2),(2012),171-177.
- [12] L.Xie and T.Xie, *Approximation theorems for localized Szász -Mirakyan operator*, J. Approx. Theory, 152(2008), 125-134.



## حول قطع مؤثر بيتا الاعتيادي

علي جاسم محمد<sup>1</sup>، هنادي عبدالله عبد الستار<sup>2</sup> وحنين جعفر صادق<sup>3</sup>

جامعة البصرة، كلية التربية للعلوم الصرفة، قسم الرياضيات<sup>1,2</sup>

جامعة البصرة، كلية العلوم، قسم الرياضيات<sup>3</sup>

[alijasmoh@yahoo.com](mailto:alijasmoh@yahoo.com), [habdulla07@gmail.com](mailto:habdulla07@gmail.com), [haneen.mahdi@yahoo.com](mailto:haneen.mahdi@yahoo.com)

## المستخلص

في هذا البحث؛ درسنا قطع مؤثر بيتا العادي  $\beta_{n,N}(x)$  والخطأ الناتج من التقريب. وضمننا خطأ القطع موصوفاً بمقياس الاستمرارية والمعيار الوزني لفضاء الدالة المقربة.

الكلمات المفتاحية : قطع مؤثر MKZ, رتبة التقريب , مؤثر بيتا , مقياس التقارب .