

PENILE SQUAMOUS CELL CARCINOMA, CASE REPORT AND REVIEW OF LITERATURES

Issam Merdan

FICMS, CABS, Assist. Prof., Department of Surgery, Collage of Medicine, University of Basrah.

Background

Penile tumors represent a difficult diagnostic and therapeutic issue to solve, mainly because of their psychological implications. The diagnosis may be delayed because many patients tend to disregard early asymptomatic lesions. Circumcision soon after birth confers almost complete immunity against carcinoma of the penis. Later circumcision does not seem to have the same effect, and Moslems circumcised between the ages of 4 and 9 years are still liable to the disease¹.

Case presentation

A 55 years old male patient presented with painful penile lesions since 6 months, associated with painful micturition and several attacks of retention of urine. He received irregular courses of different types of medical treatment without any improvement. On examination, there were two big ulcers, one anteriorly (2x3cm) and the other one (1x2cm) was laterally placed on the glans of the penis; both were irregular in shape with everted edges. The base was tender and indurated while the floor was covered by blood stained and seropurulent discharge. The bigger ulcer was reaching the urethra and invading it (Fig.1-3). Left inguinal lymph node group were enlarged, matted together and tender. Blood investigation was normal including VDRL. Ultrasound of the abdomen was normal. FNAC of the left inguinal L.N. revealed numerous acute inflammatory cells. Incisional biopsy was taken from both ulcers and the patient was kept on antibiotics until the result of the specimen revealed feature of poorly differentiated

sequamus cell carcinoma of the penis. The L.N. enlargement slightly improved on antibiotics. Because of unavailability of radiotherapy center in our governate and because of poor financial state of the patient together with the security difficulties for the patient to go to Baghdad, a decision for partial amputation of the penis was taken together with inguinal L.N. dissection. The patient run uneventful course postoperatively. Histopathology of the L.N. specimen revealed metastatic spread. The patient received chemotherapy after surgery.

Discussion

The causes of penile SCC is unclear, factors considered to play a role in the development of the disease include:

1. HPV infections: A role of HPV infections is established but not fully understood. Genital HPV infections mostly affect middle-aged, sexually active individuals².

2. In situ carcinomas: If untreated, in situ carcinomas likely evolve to invasive carcinomas. They include Bowen

disease, erythroplasia of Queyrat, and bowenoid papulosis^{3,4}.

3. Leukoplakia, Penile lichen sclerosus, Pseudoepitheliomatous, keratotic,

micaceous balanitis and Penile horn

4. Factors resulting from the lack of circumcision: Epidemiologic data have demonstrated that penile SCC is exceedingly rare in men who are circumcised at birth. The prophylactic effect of circumcision on penile carcinoma is likely related to the lack of retained smegma that has been proven to be carcinogenic in animals. One study also demonstrated an increased risk of HPV infection in uncircumcised men (19.6%) compared with circumcised men (5.5%)⁵. Poor genital hygiene in uncircumcised males, even in the absence of phimosis, may also play a role, leading to the retention of smegma³.

5. Other factors: Occasional factors that have been reported in association with penile SCC include:

- Prolonged exposure to different chemical compounds (e.g. insecticides, fertilizers)⁶.
- Late ritual circumcision followed by herbal treatments to control bleeding is associated with extensive scarring and may result in the development of invasive tumors⁷.
- Cigarette smoking, which has been found to show a clear-cut association with penile SCC related to the nicotine intake and independent of phimosis or balanitis⁶.
- UV radiation that may enhance the development of penile SCC⁸.
- Reconstructive surgery for sex reversal⁹.

Frequency: The incidence ranging from 1% in the Western countries to 10-20% in some parts of Asia, Africa, and South America¹⁰. It is rare among some religious communities who practice circumcision, such as Jewish and

Moslem groups³. The age at onset is 20-90 years, with a peak around the sixth and seventh decades¹¹.

History: In most cases, the earliest symptoms are itching or a burning sensation under the foreskin, and ulceration of the glans or the prepuce. With time, the tumor destroys the glans and the prepuce and infiltrates the corpora cavernosa¹². It most often occurs on the glans (48%), prepuce (21%), both (9%), the coronal sulcus (6%), and the shaft (<2%)¹⁰. Lymphatic metastases first occur in the superficial or deep inguinal lymph nodes, then in the regional lymphatic of the pelvis. Distant metastases resulting from vascular dissemination (most commonly to the lungs and the liver) are rare and usually late¹³. Imaging techniques recommended for a more accurate staging include ultrasonography, CT scanning, and MRI¹⁴. Initial biopsy of the primary penile lesion is necessary to confirm the diagnosis and to assess the grade and the invasiveness of the tumor³. The staging systems currently used are the Jackson classification and the (TNM) system.

Medical Care: The treatment of penile SCC varies according to the clinical stage. It includes radiation therapy, medical therapy, and surgery, alone or in combination¹⁵.

1. Radiation therapy:

Radiation therapy with external beams or mould techniques may be an option to treat small, superficial, exophytic lesions in young individuals, allowing for the preservation of sexual function with a high cure rate. In case of relapse, salvage surgery may be required. Circumcision before therapy is recommended. Major complications, such as urethral stenosis, fistulas, and penile necrosis, may occur¹⁶.

2. Medical treatment: local and systemic chemotherapy. Early premalignant and in situ changes can be treated with topical chemotherapy (5-

FU)¹⁷. Systemic chemotherapy may be used according to the stage of the disease⁽³⁾. 3. Surgical Care: Surgical procedures consist of local excision, circumcision, glansectomy, partial penectomy, total penectomy, and demasculinization. Whereas partial penectomy is the procedure of choice for tumors at a low stage (stage I, T1-T2) that involves most of the glans or the distal third of the penis. Total penectomy is necessary when the lesion is in the proximal portion of the penis or when the tumor is at an advanced stage (stages II-III, T3-T4)¹⁸.

Lymph node dissection: Because of the significant morbidity of inguinal and pelvic node dissection and because of the high incidence of reactive nodes,

surgical dissection of the lymph nodes is necessary only if they are enlarged and do not regress after adequate antibiotic therapy¹⁸.

Prevention: Circumcision in infancy and a good standard of sexual hygiene are recognized as good prophylactic measures.

Prognosis: In the absence of inguinal metastases, patients with invasive SCC of the penis involving the glans or the distal part of the shaft who undergo adequate partial amputation have a long-term survival rate of 70-80%. Of patients with involved lymph nodes, 40-50% can be cured with lymph node dissection, whereas untreated patients usually die within 2-3 years¹⁸.

Fig.1




Fig.2





Fig.3

References

1. Agarwal P, Sharma D: Auto destruction of penis due to carcinoma. *J Eur Acad Dermatol Venereal* 2005 Mar; 19(2): 220-2[[Medline](#)].
2. Bleaker MC, Hogewoning CJ, Van Den Brule AJ, et al: Penile lesions and human papillomavirus in male sexual partners of women with cervical intraepithelial neoplasia. *J Am Acad Dermatol* 2002 Sep; 47(3): 351-7[[Medline](#)].
3. Micali G, Nascia MR, Innocenzi D, Schwartz RA: Penile cancer. *J Am Acad Dermatol* 2006 Mar; 54(3): 369-91; quiz 391-4[[Medline](#)].
4. Schellhammer PF, Jordan GH, Robey EL, Spaulding JT: Premalignant lesions and nonsquamous malignancy of the penis and carcinoma of the scrotum. *Urol Clin North Am* 1992 Feb; 19(1): 131-42[[Medline](#)].
5. Castellsague X, Bosch FX, Munoz N, et al: Male circumcision, penile human papillomavirus infection, and cervical cancer in female partners. *N Engl J Med* 2002 Apr 11; 346(15): 1105-12[[Medline](#)].
6. Tsen HF, Morgenstern H, Mack T, Peters RK: Risk factors for penile cancer: results of a population-based case-control study in Los Angeles County (United States). *Cancer Causes Control* 2001 Apr; 12(3): 267-77[[Medline](#)].
7. Seyam RM, Bissada NK, Mokhtar AA, et al: Outcome of penile cancer in circumcised men. *J Urol* 2006 Feb; 175(2): 557-61; discussion 561[[Medline](#)].
8. Stern RS, Bagheri S, Nichols K; PUVA Follow Up Study: The persistent risk of genital tumors among men treated with psoralen plus ultraviolet A (PUVA) for psoriasis. *J Am Acad Dermatol* 2002 Jul; 47(1): 33-9[[Medline](#)].
9. Harder Y, Erni D, Banic A: Squamous cell carcinoma of the penile skin in a neovagina 20 years after male-to-female reassignment. *Br J Plast Surg* 2002 Jul; 55(5): 449-51[[Medline](#)].
10. Burgers JK, Badalament RA, Drago JR: Penile cancer. Clinical presentation, diagnosis, and staging. *Urol Clin North Am* 1992 May; 19(2): 247-56[[Medline](#)].
11. Aynaud O, Asselain B, Bergeron C, et al: [Intraepithelial carcinoma and invasive carcinoma of the vulva, vagina and penis in Île-de-France. Enquete PETRI on 423 cases]. *Ann Dermatol Venereal* 2000 May; 127(5): 479-83[[Medline](#)].
12. Velazquez EF, Barreto JE, Rodriguez I, et al: Limitations in the interpretation of biopsies in patients with penile squamous cell carcinoma. *Int J Surg Pathol* 2004 Apr; 12(2): 139-46[[Medline](#)].
13. Giesler EJ, Gee JR, Reece GP, et al: Cutaneous metastases of penile squamous cell carcinoma following en bloc resection of inguinal recurrence with immediate pedicle flap reconstruction. *J Urol* 2001 Oct; 166(4): 1384-5[[Medline](#)].
14. Misra S, Chaturvedi A, Misra NC: Penile carcinoma: a challenge for the developing world. *Lancet Oncol* 2004 Apr; 5(4): 240-7[[Medline](#)].
15. McDougal WS: Advances in the treatment of carcinoma of the penis. *Urology* 2005 Nov; 66(5 Suppl): 114-7 [[Medline](#)].
16. Azrif M, Logue JP, Swindell R, et al: External-beam radiotherapy in T1-2 N0 penile carcinoma. *Clin Oncol (R Coll Radiol)* 2006 May; 18(4): 320-5[[Medline](#)].
17. Klein E: Tumors of skin. IX. Local cytostatic therapy of cutaneous and mucosal premalignant and malignant lesions. *N Y State J Med* 1968 Apr 1; 68(7): 886-9[[Medline](#)].
18. Das S: Penile amputations for the management of primary carcinoma of the penis. *Urol Clin North Am* 1992 May; 19 (2): 277-82[[Medline](#)].