

Survival and metabolic activity of *lux*-marked *Escherichia coli* O157:H7 in different types of milk

Nawfal A Alhelfi¹, Rabya A Lahmer², Davey L Jones² and A Prysor Williams^{2*}

¹Department of Food Science, Agriculture College, University of Basra, Basra, Iraq

²School of Environment, Natural Resources and Geography, College of Natural Sciences, Bangor University, Bangor LL57 2UW, UK

Received 13 October 2011; accepted for publication 31 January 2012; first published online 4 April 2012

Escherichia coli O157:H7 is a potentially lethal pathogen which has been responsible for several outbreaks of milk-borne illness in recent years. The objective of this study was to evaluate the survival and metabolic activity (indexed by bioluminescence) of a chromosomally *lux*-marked strain of *Esch. coli* O157:H7 in raw, pasteurized and microfiltered pasteurized milk at 4 and 20 °C for up to 14 d. Results showed that the population of *Esch. coli* O157:H7 and its metabolic activity decreased in all samples during storage at 4 °C, with no significant differences in numbers observed between the different milk types; but metabolic activity was significantly higher ($P < 0.05$) in the microfiltered pasteurized milk than that in raw milk. At 20 °C, *Esch. coli* O157:H7 counts and cell activity peaked at day 2, and then declined progressively. At 20 °C, survival and metabolic activity were significantly lower in raw milk compared with pasteurized milk. We conclude that storage temperature is more important in regulating the survival of *Esch. coli* O157 in contaminated milk than its origin/pre-treatment conditions.

Keywords: Cross-contamination, dairy products, food poisoning, hygiene, microbiological quality.

Due to the high nutrient content of milk, it is an optimal medium for the growth of several microorganisms (Barbano et al. 2006). Consumption of raw milk, if not heat-treated or pasteurized, can be particularly problematic and is responsible for many disease outbreaks worldwide. Outbreaks are also associated with improperly pasteurized milk, and dairy products made from unpasteurized milk (Wang et al. 1997; Vernozy-Rozand et al. 2005).

Escherichia coli O157:H7 was first identified as a human pathogen in 1982 when outbreaks of bloody diarrhea and severe abdominal cramps occurred in the USA (Riley et al. 1983). The majority of affected individuals are children and the elderly, who can develop complications including haemorrhagic diarrhoea, haemolytic uremic syndrome, and thrombotic thrombocytopenic purpura (Griffin, 1995). Though only a small percentage of raw milk samples have been found to be *Esch. coli* O157:H7 positive (Duncan & Hackney, 1994), contamination with this pathogen has resulted in several milk-borne outbreaks of gastroenteritis (Chapman et al. 1993). Since *Esch. coli* O157 is an ordinary inhabitant of the bovine intestinal tract, the route of contamination with *Esch. coli* is through faecal contact

with feedstuffs, or during milking without strict hygiene practices (Hussein & Sakuma, 2005).

To date, the milk industry has successfully tackled issues of milk safety through various intervention strategies. Pasteurization has proved to be an effective measure in ensuring the safety of milk and dairy products. While unpasteurized raw milk can pose a public health concern, post-pasteurization contamination with *Esch. coli* O157 should also be noted. Faulty on-farm pasteurizers have also resulted in an outbreak of *Esch. coli* O157 (Goh et al. 2002). Incidentally, microbial growth has been shown to be greater in pasteurized samples of whey than its unpasteurized counterpart at a range of storage temperatures (Marek et al. 2004). Therefore, effective pasteurization and avoiding post-pasteurization cross-contamination in the fridge environment are both necessary to ensure the safety of milk and milk products (Heuvelink et al. 1998).

Although human pathogen outbreaks associated with milk are relatively rare, it is important to minimise this threat to maintain consumer confidence in dairy products and to protect the dairy industry. To date, no studies have examined the metabolic activity of *Esch. coli* O157:H7 in different types of milk during storage, an important evaluator of the pathogen's potential infectivity (Jawhara & Mordon, 2004). The aim of this study was to improve our understanding of the pathogen's behaviour in milk through studying both

*For correspondence; e-mail: prysor.williams@bangor.ac.uk

its survival and metabolic activity in raw and different types of pasteurized milk under ambient (20 °C) and refrigeration (4 °C) conditions. To do this, a bioluminescent (*lux*-marked) strain of *Esch. coli* O157:H7 (strain 3704 Tn5 *lux*CDABE; Ritchie et al. 2003) was used. Measuring bioluminescence in relative light units (RLU) indicates the degree of cellular metabolic activity (Ritchie et al. 2003), and has proved to be useful in improving our understanding of the pathogen in a range of contrasting environments (e.g. Jawhara & Mordon, 2004; Williams et al. 2008a, b; Thorn et al. 2011).

Materials and Methods

Preparation of milk

Raw milk was collected from the tank of a dairy farm located in Bangor, North Wales. The samples were kept at 4 °C in sterile ice bags during transportation. Milk was used within 3 h after arrival at the laboratory. Part of the raw milk remained unpasteurized, whilst part was heat-treated in glass containers to 63.5 °C (30 min) to prepare laboratory-pasteurized milk. Fresh full-fat commercially-pasteurized and full-fat microfiltered pasteurized milk (Cravendale) were purchased from Arla Foods UK Ltd (Leeds, UK).

Screening milk samples for *Esch. coli* O157:H7

Milk samples were tested for the presence of *Esch. coli* O157 before inoculation. Isolation and detection of *Esch. coli* O157 involved enrichment followed by immunomagnetic separation (IMS). To start with, 5 ml of each milk samples were mixed with 45 ml of modified Tryptone Soy Broth (mTSB) (Oxoid CM 0989; Oxoid Ltd., Basingstoke, UK) and incubated at 37 °C for 6 h. Afterwards, 1 ml of the enriched sample was analysed by DynamagTM-2 IMS (Invitrogen Dynal A.S., Oslo, Norway) with 0.02 ml of Captivate[®] *Esch. coli* O157 immunomagnetic beads (Lab M Ltd, Bury, UK) and incubated at 25 °C for 30 min. After IMS, the beads were washed three times using phosphate buffered saline with 0.05% Tween 20 as wash buffer, and resuspended in 0.1 ml of the same buffer. They were then spread equally on three SMAC plates (sorbitol MacConkey agar plates (SMAC; Oxoid CM813) supplemented with cefixime (0.05 mg/l) and potassium telluride (2.5 mg/l) CT-SMAC), and incubated at 37 °C for 18 to 24 h.

Inoculation of milk samples with *Esch. coli* O157:H7

An inoculum was prepared from a fresh overnight culture (LB broth; Difco Ltd, Teddington, Surrey, UK; 18 h, 37 °C, 150 rev./min) of *Esch. coli* O157:H7 (Ritchie et al. 2003) in stationary growth phase. Cells were washed and concentrated by centrifugation as described in Avery et al. (2005). An inoculum (1 ml) of the mixture at the appropriate dilution was added to 99 ml of each milk type and mixed thoroughly in sterilised screw-cap bottles to obtain the desired final concentration of approximately 10³ CFU/ml. All bottles of

inoculated milk and uninoculated milk (control) were incubated at 4 and 20 °C.

Survival and metabolic activity of *Esch. coli* O157:H7

Esch. coli O157 cells were enumerated at 0 (immediately after inoculation), 1, 2, 4, 6, 8, 10, 12 and 14 d post-inoculation. Milk samples were serially diluted in Ringer solution (Oxoid), and serial dilutions were plated onto CT-SMAC and incubated at 37 °C for 18 to 24 h. Non-sorbitol fermenting *Esch. coli* O157:H7 colonies were confirmed by agglutination with a latex test kit (Oxoid DR0620).

A parallel experiment was designed to assess variations in the activity of *Esch. coli* O157 among the different milk types (raw, laboratory-pasteurized, full-fat commercially-pasteurized, and microfiltered pasteurized). Bioluminescence of bacteria in milk was measured at 0 (immediately after inoculation), 1, 2, 4, 6, 8, 10, 12 and 14 d post-inoculation. At each time-point, a 1-ml aliquot from samples used for the enumeration study detailed above was placed into a plastic luminometer cuvette and its luminescence (RLU) was determined using a SystemSURE plus Pi-102 Luminometer (Hygiena International Ltd, UK).

Aerobic plate counts and pH

Aerobic plate counts (APC) were determined from uninoculated milk samples (control) at 0 (immediately after inoculation), 1, 2, 4, 6, 8, 10, 12 and 14 d. The uninoculated samples were serially diluted in Ringer solution, and serial dilutions (1:10) were plated onto plate count agar (PCA; Oxoid) and incubated at 30 °C for 48 h.

Samples' pH values were determined with a standard pH meter (Hanna instruments pH 211). Calibration was performed using two standard buffer solutions at pH 4.0 and 7.0.

Statistical analysis

Outcomes in the experiment were changes in *Esch. coli* O157:H7 cell counts and cell activity (bioluminescence), aerobic plate counts, and pH values during the 14 d incubation period. Log ($y+1$) transformation was performed on *Esch. coli* O157:H7 cell counts and cell activity, aerobic plate counts, which together with pH data were subjected to ANOVA tests and Tukey's test with significance at $P<0.05$ using SPSS 18.0 software (SPSS Inc, Chicago, Illinois, USA).

Results

Screening milk samples for *Esch. coli* O157:H7

No *Esch. coli* O157:H7 was detected by the IMS method in any of the milk samples before inoculation.




Fig. 1. Survival and metabolic activity of *Esch. coli* O157:H7 in milk samples stored at 4 and 20 °C (R=raw milk, LP=laboratory-pasteurized milk, F=microfiltered pasteurized milk and P=commercially-pasteurized milk). Values represent means \pm SEM ($n=3$).

Survival and metabolic activity of *Esch. coli* O157:H7

Survival and metabolic activity of *Esch. coli* O157:H7 at both 4 and 20 °C are shown in Fig. 1. At 4 °C, *Esch. coli* O157:H7 populations declined steadily and continuously by 1.0–1.5 log₁₀ CFU/ml in all samples over 14 d incubation. While log cell count reduction was greatest in raw milk (1.5 log₁₀ CFU/ml), between-sample variations in survival of *Esch. coli* O157:H7 were not significant between all samples at this temperature ($P>0.05$). Metabolic activity of *Esch. coli* O157:H7 continuously and steadily reduced (by 1.3–2.07 log₁₀ RLU) over the 14 d, with activity in raw milk diminishing near to zero. Cell activity in the microfiltered milk was significantly higher than that in raw milk ($P<0.05$), while no significant difference was seen among laboratory-pasteurized, commercially-pasteurized and raw milk ($P>0.05$).

At 20 °C, *Esch. coli* O157:H7 cell counts in all milk samples showed a dramatic initial increase, peaking at day 2 (2.7–3.6 log₁₀ CFU/ml), then progressively declined until the end of the 14 d incubation. Cell counts in raw milk samples decreased most (about 2.5-log cell count reduction using day 0 as baseline) and the count reduction was significantly higher ($P<0.001$) in raw milk compared with pasteurized samples. Counts did not statistically differ between the different types of pasteurized milk. Further ANOVA tests revealed that temperature was a significant factor moderating survival

in all samples, with higher environmental temperatures leading to higher pathogen counts ($P<0.001$). Metabolic activity of *Esch. coli* O157:H7 at 20 °C in all milk samples increased significantly on day 1, which continued to rise and peak (2.3–2.75 log₁₀ RLU) at day 2. Cell activity dropped significantly afterwards in all samples, reaching zero in raw milk at day 6 and at day 10 in pasteurized milk samples (Fig. 1). Cell activity in the three pasteurized milk samples was significantly higher than that in raw milk ($P<0.05$). Further ANOVA analysis indicated that temperature was an important influence on *Esch. coli* O157:H7 cell activity, with the higher temperature inducing a peak at day 2 which was not observed at the low temperature.

Aerobic plate counts and pH

Changes in APC (measured in log₁₀ CFU/ml) and pH values are shown in Fig. 2. At 4 °C, average APC increased significantly more in raw (around 3.9 log count growth) than in pasteurized milk (<1.35 log count growth) (all $P<0.001$, mean counts: raw>laboratory-pasteurized>commercially-pasteurized>microfiltered). At 20 °C, APC increased approximately 5–6 log₁₀ CFU/ml in all milk samples after 14 d of incubation. APC in raw milk were significantly higher ($P<0.001$) than that in the three types of pasteurized milk,




Fig. 2. Changes in aerobic plate counts (measured in \log_{10} CFU/ml) and pH in milk samples stored at 4 and 20 °C (R=raw milk, LP=laboratory-pasteurized milk, F=microfiltered pasteurized milk and P=commercially-pasteurized milk). Values represent means \pm SEM ($n=3$).

with values in microfiltered milk being significantly lower than those in the other pasteurized milk types ($P<0.001$). In the ANOVA test, temperature was found to be a significant factor in aerobic cell growth in the pasteurized milk samples. The growth rates in pasteurized samples were significantly higher under room temperature than under refrigeration temperature; although growth slowed down from day 8. In all, there was a final increase of around 6 \log_{10} CFU/mL in APC from day 0 to 14.

At 4 °C, no substantial changes in pH were observed in the pasteurized milk samples (Fig. 2), staying between 6.5 and 6.7; however pH values in raw milk exhibited a gradual decrease to 5.7. Over 14 d at 20 °C, pH values decreased rapidly in all samples, from an average of 6.7 to 3.3 in raw milk, and to around 4.0–4.2 in pasteurized milk samples. Over the course of the experiment, there were significant differences in pH among the four types of milk (all $P<0.001$, pH mean: microfiltered > commercially-pasteurized > laboratory-pasteurized > raw). Changes in pH values were negatively associated with the increase in APC, with higher numbers of aerobic microorganisms leading to lower pH values.

Discussion

Pasteurized and unpasteurized milk may be contaminated with *Esch. coli* O157:H7 when inadequate farm hygiene measures (milking and milk handling) are present or post-pasteurization contamination occurs. Given the low infective dose of *Esch. coli* O157:H7 (Chart, 2000) and the association of milk with past infections, it is important to understand the behaviour of the organism in dairy products. Whilst others have previously studied changes in numbers of the organism in dairy products (e.g. Wang et al. 1997; Mamani et al. 2003; Marek et al. 2004), this is the first study to concurrently monitor the pathogen's metabolic activity. Given the association between metabolic activity and infectivity, this paper presents novel findings of interest to dairy microbiology and food safety.

The present study confirmed that temperature is an important factor that influences the survival and activity of *Esch. coli* O157:H7. We observed that *Esch. coli* O157:H7 could not grow under refrigeration conditions in any type of milk, which was largely consistent with results from previous studies on a limited range of milk types (Wang et al. 1997).

Previous studies have recommended that milk be kept at $\leq 5^{\circ}\text{C}$ as even at 7°C , *Esch. coli* O157 can grow at a significant rate (Heuvelink et al. 1998). Whilst other studies have also found the organism to survive and proliferate at room temperatures (Wang et al. 1997; Mamani et al. 2003), this study additionally revealed a corresponding increase in the pathogen's metabolic activity at elevated temperatures.

Esch. coli O157:H7 numbers and metabolic activity consistently decreased at a greater rate in raw milk than in the three types of pasteurized milk. Greater APC values were recovered from raw milk and this is expected to result in elevated competition with, and/or antagonism against the pathogen, as reported elsewhere (Wang et al. 1997; Elwell & Barbano, 2006). Storage of raw milk at 20°C also reduced pH considerably, most probably due to lactic acid production by the elevated counts of background microorganisms (Kuipers et al. 2000). Acidic conditions ($\text{pH} < 3.5$, Fig. 2) are likely to be detrimental to survival of *Esch. coli* O157:H7; however it should be noted that the pathogen was not found to be eliminated at these low pH values, consistent with previous studies that show its acid resistance and adaptation in acidic environments (Leyer et al. 1995; Mamani et al. 2003; Carter et al. 2011). In addition, raw milk may also contain several compounds with bioactive components (e.g. lactoferrin, lactoperoxidase and lysozyme) that can reduce or eliminate populations of pathogenic bacteria; however these will be lost during heat treatment (IDF, 1991).

To conclude, examining the role of incubation temperature has practical significance in understanding how *Esch. coli* O157:H7 and other aerobic cells behave in the food chain, from retailer fridge storage to the consumer home where the greatest risk of human infection occurs. We have shown that allowing contaminated milk to reach room temperature for even a space of 2 h can induce a transient proliferation of *Esch. coli* O157:H7 numbers and metabolic activity. Microfiltering milk did not have a significant effect on pathogen proliferation in comparison to normal pasteurization procedures and especially in comparison to the importance of storage temperature. Although pasteurization represents an effective measure to reduce pathogenic risks and improves the microbial quality of milk, consistent hygiene quality standards must be observed both pre- and post-pasteurization to guard against any possible pathogen and spoilage microorganisms.

This work was undertaken using funds provided by the UK Rural Economy and Land Use Programme (RES-229-25-0012).

References

- Avery LM, Killham K & Jones DL 2005 Survival of *Esch. coli* O157:H7 in organic wastes destined for land application. *Journal of Applied Microbiology* **98** 814–822
- Barbano DM, Ma Y & Santos MV 2006 Influence of raw milk quality on fluid milk shelf life. *Journal of Dairy Science* **89** E15–E19
- Carter MQ, Brandl MT, Louie JW, Kyle JL, Carychao DK, Cooley MB, Parker CT, Bates AH & Mandrell RE 2011 Distinct acid resistance and survival fitness displayed by curli variants of enterohemorrhagic *Escherichia coli* O157:H7. *Applied and Environmental Microbiology* **77** 3685–3695
- Chapman PA, Wright DJ & Higgins R 1993 Untreated milk as a source of verotoxigenic *Escherichia coli* O157. *Veterinary Record* **133** 171–172
- Chart H 2000 VTEC enteropathogenicity. *Journal of Applied Microbiology Symposium Supplement* **88** 12S–23S
- Duncan SE & Hackney CR 1994 Relevance of *Escherichia coli* O157:H7 to the dairy industry. *Dairy, Food and Environmental Sanitation* **14** 656–660
- Elwell MW & Barbano DM 2006 Use of microfiltration to improve fluid milk quality. *Journal of Dairy Science* **89** 10–30
- Goh S, Newman C, Knowles M, Bolton FJ, Hollyoak V & Richards S 2002 *Esch. coli* O157 phage type 21/28 outbreak in North Cumbria associated with pasteurized milk. *Epidemiology and Infection* **129** 451–457
- Griffin PM 1995 *Escherichia coli* O157:H7 and other enterohemorrhagic *Escherichia coli*. In *Infections of the Gastrointestinal Tract*, pp. 739–762 (Eds MJ Blaser, PD Smith, JI Ravdin, HB Greenberg & RL Guerrant). New York: Raven Press, Ltd
- Heuvelink AE, Bleumink B, van den Biggelaar FLAM, Te Giffel MC, Beumer RR & de Boer E 1998 Occurrence and survival of verocytotoxin-producing *Escherichia coli* O157 in raw cow's milk in the Netherlands. *Journal of Food Protection* **61** 1597–1601
- Hussein HS & Sakuma T 2005 Prevalence of Shiga toxin producing *Escherichia coli* in dairy cattle and their products. *Journal of Dairy Science* **88** 450–465
- International Dairy Federation (IDF) 1991 *Detection and Confirmation of Inhibitors in Milk and Milk Products*. IDF – Bulletin No. 258. Brussels, Belgium: IDF
- Jawhara S & Mordon S 2004 *In-vivo* imaging of bioluminescent *Escherichia coli* in a cutaneous wound infection model for evaluation of an antibiotic therapy. *Antimicrobial Agents and Chemotherapy* **48** 3436–3441
- Kuipers OP, Buist G & Kok J 2000 Current strategies for improving food bacteria. *Research in Microbiology* **151** 815–822
- Leyer GJ, Wang LL & Johnson EA 1995 Acid adaptation of *Escherichia coli* O157:H7 increases survival in acidic foods. *Applied and Environmental Microbiology* **61** 3752–3755
- Mamani Y, Quinto EJ, Simal-Gandara J & Mora MT 2003 Growth and survival of *Escherichia coli* O157:H7 in different types of milk stored at 4°C or 20°C . *Food Microbiology and Safety* **68** 2558–2563
- Marek P, Nair MKM, Hoagland T & Venkatarayanan K 2004 Survival and growth characteristics of *Escherichia coli* O157:H7 in pasteurized and unpasteurized Cheddar cheese whey. *International Journal of Food Microbiology* **94** 1–7
- Riley LW, Remis RS, Helgeson SD, McGee HB, Wells JG, Davis BR, Hebert RJ, Olcott ES, Johnson LM, Hargrett NT, Blake PA & Cohen ML 1983 Hemorrhagic colitis associated with a rare *Escherichia coli* serotype. *New England Journal of Medicine* **308** 681–685
- Ritchie JM, Campbell GR, Shepherd J, Beaton Y, Jones D, Killham K & Artz RRE 2003 A stable bioluminescent construct of *Escherichia coli* O157:H7 for hazard assessments of long-term survival in the environment. *Applied and Environmental Microbiology* **69** 3359–3367
- Thorn CE, Quilliam RS, Williams AP, Malham SK, Cooper D, Reynolds B & Jones DL 2011 Grazing intensity is a poor indicator of waterborne *Escherichia coli* O157 activity. *Anaerobe* **17** 330–333
- Vernozy-Rozand C, Montet MP, Berardin M, Bavai C & Beutin L 2005 Isolation and characterization of Shiga toxin-producing *Escherichia coli* strains from raw milk cheeses in France. *Letters in Applied Microbiology* **41** 235–241
- Wang GD, Zhao T & Doyle MP 1997 Survival and growth of *Escherichia coli* O157:H7 in unpasteurized and pasteurized milk. *Journal of Food Protection* **60** 610–613
- Williams AP, Gordon H, Jones DL, Strachan NJC, Avery LM & Killham K 2008a Leaching of bioluminescent *Escherichia coli* O157:H7 from sheep and cattle faeces during simulated rainstorm events. *Journal of Applied Microbiology* **105** 1452–1460
- Williams AP, McGregor KA, Killham K & Jones DL 2008b Persistence and metabolic activity of *Escherichia coli* O157:H7 in farm animal faeces. *FEMS Microbiology Letters* **287** 168–173