

EFFECT MICROENCAPSULATION OF *Saccharomyces boulardii* ON VIABILITY OF YEAST IN VITRO AND ICE CREAM

Alaa Kareem Niamah^{*1}, Alaa Jabbar Al-Manhel¹, Shayma Thyab Gddoa Al-Sahlany¹

¹ Department of Food Science, College of Agriculture, University of Basrah, Basrah City, Iraq
^{*}alaakareem2002@hotmail.com

Article history:

Received:

26 July 2017

Accepted:

1 September 2018

Keywords:

Saccharomyces boulardii;

Microencapsulation;

Probiotic;

Ice cream.

ABSTRACT

The effect of microencapsulation by alginate and alginate-chitosan on the survival of *Saccharomyces boulardii* ATCC MYA-796 in simulated gastric and intestinal juices was investigated. The survival of yeast microencapsulated cells treated in ice cream production. The percentage survival of alginate and alginate-chitosan encapsulated cells treated in simulated gastric juice after 240 was 80 % and 90 % respectively, simulated intestinal juice was 80% and 85% respectively. The viability of *Saccharomyces boulardii* (Log CUF/g) in ice cream after storage time (at freezing temperatures 21 day) was 7.11, 6 and 5.25 for alginate-chitosan microcapsules treatment, alginate treatment and free cell treatment respectively. The pH values range of ice cream in three treatments at the initial storage time was 4.70-4.77. After 21 days, the pH decreased in all treatments.

1.Introduction

Probiotics have been defined as live microbes, which transit the gastro-intestinal tract, and in doing so benefit the health of the consumer. Generally, probiotic microorganisms have a therapeutic effect for improved immune system (by microbial metabolites, cell wall components and DNA) (FAO/WHO, 2002). As well as the treatment of diarrhoea (antitoxin effects) by *Saccharomyces boulardii* and *Saccharomyces cerevisiae* are the most common yeast strains that have desirable properties used in probiotic products (Nousia et al., 2011; Arslan et al., 2016). *S. boulardii* is unique probiotic and bio-therapeutic yeast, known to survive in gastric acidity and it is not adversely affected or inhibited by antibiotics or does not alter or adversely affect the normal microflora in the bowl. *S. boulardii* has been utilized worldwide as a probiotic supplement to

support gastrointestinal health. In recent years, by incorporating *S. boulardii* various dairy foods such as yoghurt, ultra high temperature treated (UHT) milk, acidophilus yeast milk, ice cream etc. (Hattingh, and Viljoen, 2001; Karaolis et al., 2013, Niamah, 2017). Ice cream is a product with peculiar textural and organoleptic features and is highly appreciated by a very broad spectrum of consumers. Ice cream's structure and colloidal design, together with its low-temperature storage, renders it a very promising carrier for the stabilization and in vivo delivery of bioactive compounds and beneficial microorganisms. (Soukoulis et al., 2014), However, temperature change, also referred to as "cold shock", during freezing and melting may cause damage such as reduction or even complete loss of metabolic activity (Mohammadi et al., 2011). A method for

encapsulation and stabilization of probiotic yeast as *S. boulardii* or bacterial strains (*Lactobacillus* spp. and *Bifidobacterium* spp.) in polymeric or biopolymeric fibers has been developed, biopolymers are natural polymers (chitosan, alginate) that are abundantly available and extractable from natural sources and these biopolymers offer a wide range of unique applications (Abd and Niamah, 2012; Al-Manhel et al. 2018). In order to achieve the claimed beneficial effects of probiotic bacteria, these specific microorganisms must be viable, active, and abundant in the product up to the expiry date (cell counts range from 10^6 to 10^9 CFU/g) (Casarotti and Penna, 2015). Microencapsulation is a process by which bioactive materials are coated with other protective materials or their mixtures (Huq et al., 2013). Ice cream has nutritional significance, but possesses no therapeutic properties. The growing interest of consumers in therapeutic products has led to the incorporation of probiotic cultures into ice cream to result in a dietetic ice cream. Some studies have demonstrated that it is possible to produce probiotic ice cream using different microorganism (Ahmadi et al., 2012). This study was undertaken to select a suitable for *S. boulardii* microencapsulation and checked for their viability, survival of the encapsulated probiotics under the stimulated gastric conditions and study was to manufacture a probiotic ice cream containing *S. boulardii* and to determine how long these yeasts would remain viable during frozen storage of the ice cream.

2. Materials and methods

2.1. Materials

2.1.1. Reagents

Sodium alginate and chitosan were purchased from (BDH, UK). Whole cow milk obtained from Animals station of Agriculture College / University of Basrah.

2.1.2. Yeast cell culture

S. boulardii ATCC MYA-796™ was obtained from Department of Food Science /

College of Agriculture university of Basrah and grow in YPD media (content: 0.3% yeast extract, 0.5% peptone and 1% glucose) at 30°C for 48 h.

2.2. Methods

2.2.1. Yeast inoculum preparation

33×10^9 CFU/ mL of *S. boulardii* was inoculated in 50 ml of YPD media and incubated at 30°C at 150 rpm in a shaking incubator for 48 h. The cells were harvested by spinning them at 5,000 rpm for 10 min. The cultures were then washed twice with sterile saline solution (0.9%) and used in the microencapsulation process.

2.2.2. Microencapsulation of yeast cells

A 3.0% of alginate solution was autoclave sterilized at 121°C for 20 min and stored at 4°C. After washing, the cells were suspended in 5 mL of sterile distilled water and mixed with 20 mL of 2% (w/v) sodium alginate for at least 30 min. The cell suspension was placed in a sterile syringe and injected through a 0.11 mm needle into sterile 0.05 M CaCl_2 . The 0.11 mm needle was used to produce beads with a diameter of 1-2 mm. Moreover, stirring with a magnetic stirrer was applied during dropping to produce spherical beads, and the distance between the needle and the surface of the solution was controlled to approximately 1 to 2 cm to avoid formation of flat beads. After 30 min gelification in CaCl_2 , the beads were rinsed with, and then kept in, sterile water at 4°C (Krasaekoopt et al., 2004).

2.2.3. Preparation of alginate-chitosan microcapsules entrapping yeast cells

Chitosan solution (0.5% w/v) was prepared by the addition of chitosan to acetic acid solution (0.1M) with stirring until dissolution (~1 h). The calcium alginate microcapsules were immersed in 100 ml of chitosan solution and shaken at 100 rpm for 40 min on an orbital shaker for coating and then filtration. The chitosan-coated beads were washed and kept in sterile water at 4°C (Krasaekoopt et al., 2004).

2.2.4. Preparation of simulated gastric and intestinal juices and inoculation of cells

The simulated juices were prepared according to Brinques et al. (2011). Simulated gastric juices were prepared by dissolving pepsin (Riedel-DeHaen Hannover) in sterile sodium chloride solution (0.5% w/v) to a final concentration of 3.0 g/L and adjusting the pH to 1.5 with hydrochloric acid. Simulated intestinal juices were prepared by suspending pancreatin (BDH) in sterile sodium chloride solution (0.5%, w/v) to a final concentration of 1 g/L with 4.5% bile salts (Oxoid, UK) and adjusting the pH to 8.0 with sterile NaOH (0.1 M). Both solutions were filtered for sterilization through a 0.22 µm membrane. *S. boulardii* inoculated to the simulated gastro-intestinal juice individually in three different forms, non-encapsulated, encapsulated with calcium alginate and encapsulated with alginate and chitosan. Then one gram of freshly encapsulated yeast samples or 1 mL of cell suspensions (free cells) were gently mixed with 10 mL of sterile simulated gastric juice or sterile simulated intestinal juice and incubated at 37 °C for 60, 120, 180 and 240 min. Surviving yeast were enumerated by pour plate counts in YPG agar aerobically incubated at 30°C for 48 h. the survival percentage of yeast was calculated by the following formula:

$$\text{Survival of yeast cells \%} = \frac{\text{Final Log (CFU/mL)}}{\text{Control Log (CFU/mL)}} \times 100 \quad (1)$$

2.2.5. Preparation of ice cream

Ice cream mix was prepared by mixing ingredients and then heated to 80°C for 30 sec. Mixes were cooled to 5°C and aged overnight at the same temperature. After ageing, the mix was heat treated to 80°C for 30 sec. and cooled to 37°C. Yeast probiotic (*S. boulardii*) were inoculated into ice cream mix at the rate of 5% (33×10^9 CFU/ mL) and incubated at 30°C for 5 h. The ice cream was filled in 50 ml food grade paper cups, covered with food grade lids and stored at -18°C (Pandiyan, 2010). The probiotic ice cream was prepared using the different treatments as shown below: Treatment I: Ice

cream prepared with free cell of *S. boulardii*
Treatment II : Ice cream prepared with sodium alginate microcapsules entrapping yeast cells.
Treatment III: Ice cream prepared with alginate-chitosan microcapsules entrapping yeast cells.

2.2.6. Yeast viability

Frozen fermented ice cream was thawed and then diluted. The viability of free and microencapsulated yeast cells was enumerated by pour plate count method. To determine the probiotic viability count, the entrapped probiotics were released from the microcapsules. One gram of the ice cream samples transfers to 9 mL of sterile peptone water and then homogenized for 15sec. in a vortex mixer. The success of the encapsulating formula was tested. Thus, one-gram sample of microcapsules was diluted in sterile peptone water and then an aliquot of 100 mL was discharged into potato dextrose agar plates and incubated at 30°C for 24–72 h. The population of *S. boulardii* was then quantified and expressed as the number of colony-forming units (CFU) per gram (CFU/g) Samples were taken at 0, 1, 7, 14 and 21 days of storage (Zamora-vega et al., 2012). The pH of the samples was measured with a pH meter (SD-300, Germany) (Niamah et al., 2016).

3. Results and discussions

3.1. The survival *S. boulardii* in simulated gastric juice

The survival of the free cells and encapsulated *S. boulardii* in simulated gastric juice after being kept for 240 min is shown in Figure 1. Food stays in the stomach for 2 to 4 hours, while the liquids can only stay about 20 minutes. pH of gastric juice simulates affected the survival of both the free and encapsulated cells of *S. boulardii*. The encapsulated *S. boulardii* survived well in simulated gastric conditions compared to the free yeast cells and the encapsulated by alginate-chitosan was best than with alginate. The percentage survival of encapsulated *S. boulardii* in gastric juice after 3 h was 89% and 92%, for alginate

microcapsules method and alginate-chitosan microcapsules method while the percentage was 80% free yeast cells. After 240 min. the difference between alginate and alginate-chitosan microcapsules was more clearly in results. Whereas the yeast in the alginate-only microcapsules loss of percentage survival to 82%, the chitosan-mixed alginate microcapsules was 90%.

Alginate mixture remains stable structurally in acidity environments when the pH decreases as the viscosity of alginate solutions increase

and the carboxylate groups in the alginate solutions backbone become protonated lead to form hydrogen bonds, which increased bindings yeast cells (Lee and Mooney, 2012). Encapsulation, either by alginate or by alginate/chitosan provide the best protection in the case of yeast cells encapsulated compared with non-encapsulated cells (free cells). This can be attributed to the polysaccharides of chitosan acting as a buffer, reducing the activity of the acid. It does protective effect to yeast cells.




Figure 1. The percentage survival of yeast in simulated gastric

3.2.The survival *S. boulardii* in simulated intestinal juice

Bile salt tolerance is one of the basic characteristics of the probiotic to survive in the small intestine. Survival of encapsulated and free cells of *S. boulardii* after exposure to 4.5% bile salt for 240 min. This reflects the time of the stay of food in the small intestine is shown in Figure 2. The percentage survival of encapsulated *S. boulardii* was 80% and 85% after 240 min for alginate microcapsules method and alginate-chitosan microcapsules method while percentage survival of *S. boulardii* free cells was 67%. It was noted that the survival of alginate encapsulated and alginate-chitosan encapsulated *S. boulardii* reduced to somewhat compared with *S. boulardii* free cells. Mixed the alginate microcapsulation with chitosan more improve the viability of *S. boulardii* (Cook et al., 2011).

The results of the study agreed with the (Dikit et al, 2015) who studied the effect of encapsulation on the survival of *Lactobacillus plantarum*. pH-base in the small intestine affecting the survival of *S. boulardii* cells. Microcapsulation of *S. boulardii* cells does to prevent the effect of the acid and bile salts. Chitosan polysaccharide chitosan is positively charged, and therefore, it is not suitable for encapsulated yeast when it is used by itself. A mixture of chitosan with other biological substances usually soluble in solvents such as alginate. Alginate is a negatively charged polysaccharide and best substance for growth and microcapsulation of probiotic microorganisms. Crosslinked three-dimensional made of chitosan and alginate provides much

better for microcapsulation process (Baysal et al., 2013).




Figure 2. The percentage survival of yeast in intestinal environment

3.3. The viability of *S. boulardii* in ice cream

S. boulardii grew to high numbers in ice cream mix. Even the high solids level of the ice cream mix did not prevent growth of either free cells or microcapsule when a high percentage (4%) of inoculum was used. Viable cell count of *S. boulardii* (free cells) decreased from 8.55 to 5.25 log CFU/g due to encapsulation process; Changes in viable counts of *S. boulardii* in ice cream mix due to freezing and during frozen storage at -18°C are presented in Figure 3. Results showed that there was an approximately 3.3 Log cycle decrease in the count of probiotic cells in the free form immediately after freezing for 3 week compared with the encapsulated probiotic cells decreased 2.51 and 1.51 Log cycle in Treatment II and Treatment III respectively. This indicates that freezing had destructive effects on probiotic cells most probably due to the freezing injury of cells. During the freezing process, the cells of probiotics can be lethally injured by damage to their cell walls or membranes caused by thawing and the mechanical stresses (generated by mixing and incorporating oxygen into the mixture, during the manufacture of ice creams) of ice crystals forming in the external medium or inside the cells, by cold injuries and temperature decrease shock to the cells, by condensation of solutes in the extracellular/intracellular medium, or by

dehydration of the cells. All these factors may result in even lower viability (Akin et al. 2007; Mohammadi et al., 2011). According to several findings, encapsulation is a useful alternative to increase the survival rate of probiotic bacteria in ice cream and fermented frozen dairy desserts. This result agrees with Ahmadi et al. (2012) has found that the viable counts of free probiotics decreased from ~ 9.55 to ~ 7.3 Log CFU/g after 60 days of frozen storage while that of encapsulated cells merely decreased less than 1 Log cycle. Encapsulation with alginate microbeads protected the probiotic cells against injuries in the freezing stage as well as, during frozen storage. Homayouni et al. (2008) reported that the survival of *L. casei* and *B. lactis* were monitored during the product's storage for 180 days at -20°C . The viable cell number of *L. casei* and *B. lactis* in the free state in prepared ice cream mixture was 5.1×10^9 and 4.1×10^9 CFU/mL at day one and after 180 days storage at -20°C , these numbers were decreased to 4.2×10^6 and 1.1×10^7 CFU/mL respectively. When encapsulated the mentioned probiotic bacteria in calcium alginate beads, the probiotic survival raised at a rate of 30% during the same period of storage at the same temperature. Akin et al. (2007) and Heydari et al. (2012) reported a similar observation. Encapsulation within alginate microbeads and alginate-chitosan protected the probiotic cells

against injuries in the freezing stage (Figure 3) as well as, during frozen storage. Viable number of encapsulated *L. acidophilus* was

merely decreased less than 1 Log cycle after 60 days of frozen storage.




Figure 3. Viability of free and encapsulated *S. boulardii* (Log CFU/g) in ice cream during storage periods at freezing temperature.

3.4. pH of ice cream

Table 1. Show that the pH values of three ice cream treatments after process was between 4.70-4.77. The values decreased after storage for up to 4.20 in alginate-chitosan microcapsules treatment, 4.45 in alginate microcapsules treatment and 4.45 in free cell treatment. The reduced pH because of the ability of yeast to ferment sugars in the ice-cream mixture. *S. boulardii* either by acid hydrolysis or by producing enzymes that cleave

the sucrose into glucose and fructose and glucose utilization rate was much faster than the fructose and sucrose (Kurtzman et al., 2011). *S. boulardii* is acetate produce from sugar utilization and another carbohydrates (Anjum et al., 2010). The high viability cells of *S. boulardii* in alginate-chitosan microcapsules treatment led to increased acidity and reduced pH compared with other treatments.

Table. 3 pH of ice cream treatment

Cell	pH				
	Time (day)				
	0	1	7	14	21
Alginate microcapsules	4.77	4.72	4.62	4.53	4.45
	± 0.03	± 0.04	± 0.01	± 0.02	± 0.01
Alginate-chitosan microcapsules	4.70	4.65	4.55	4.43	4.20
	± 0.03	± 0.01	± 0.02	± 0.00	± 0.05
Free cell	4.75	4.70	4.63	4.60	4.54
	± 0.05	± 0.01	± 0.01	± 0.10	± 0.06

±: Standard division (SD)

4. Conclusions

Microcapsulation during the emulsion technology was an effective way to increase the survival of probiotics yeasts as they pass during

digestive tract conditions of warm-blooded animals. Adding chitosan when using encapsulation techniques can improve the survival of *S. boulardii* in simulated gastric

juice and intestinal juice. Probiotic ice cream can be manufactured using encapsulated *S. boulardii* to ferment ice cream mix. Coating alginate microcapsules with chitosan further improve the viability of *S. boulardii* during storage time of ice cream production. The probiotic yeast *S. boulardii* can be well incorporated into ice cream based foods to develop functional and therapeutic foods.

5. References

- Abd, A.J., Niamah, A.K. (2012). Effect of chitosan on apple juice quality. *International Journal of Agriculture and Food Science*, 2(4), 153-157.
- Ahmadi, A., Milani, E., Madadlou, A., Mortazavi, S.A., Mokarram, R.R. , Salarbashi, D. (2014). Synbiotic yogurt-ice cream produced via incorporation of microencapsulated *Lactobacillus acidophilus* (LA-5) and fructooligosaccharide. *Journal of Food Science and Technology*, 51(8), 1568–1574.
- Akin, M.B., Akin, M.S. , Kirmaci, Z. (2007). Effects of inulin and sugar levels on the viability of yogurt and probiotic bacteria and the physical and sensory characteristics in probiotic ice cream. *Food Chemistry*, 104(1), 93–99.
- Akin, M.S. (2005). Effects of inulin and different sugar levels on viability of probiotic bacteria and the physical and sensory characteristics of probiotic-fermented ice cream. *Milchwissenschaft*, 60, 297–300.
- Al-Manhel, A.J., Al-Hilphy, A.R.S. , Niamah, A.K. (2016). Extraction of chitosan, characterization and its use for water purification. *Journal of the Saudi Society of Agricultural Sciences*, 17, 186–190.
- Anjum, F. M., Sabir, M. Z., Khan, M. I., Pasha, I. (2010). Sugar utilization behavior of yeast (*Saccharomyces cerevisiae*) types and doses on bread quality. *Nutrition & Food Science*, 40(4), 395–402.
- Arslan , A.A. , Gocer, E.M.C., Demir, M., Atamer, Z., Hinrichs, J., Küçükçetin, A. (2016). Viability of *Lactobacillus acidophilus* ATCC 4356 incorporated into ice cream using three different methods. *Dairy Science & Technology*, 96(4), 477–487.
- Baysal, K., Aroguzc, A.Z., Adiguzela, Z., Baysala, B.M. (2013). Chitosan/alginate crosslinked hydrogels: Preparation, characterization and application for cell growth purposes. *International Journal of Biological Macromolecules*, 59, 342–348.
- Brinques, G.B., Ayub, M.A.Z. (2011). Effect of microencapsulation on survival of *Lactobacillus plantarum* in simulated gastrointestinal conditions, refrigeration, and yogurt. *Journal of Food Engineering*, 103, 123-128.
- Casarotti, S.N., Penna, A.L.B. (2015). Acidification profile, probiotic invitro gastrointestinal tolerance and viability in fermented milk with fruit flours. *International Dairy Journal*, 41, 1–6.
- Cook, M.T., Tzortzis, G., Charalampopoulos, D., Khutoryanskiy, V.V. (2011). Production and evaluation of dry alginate-chitosan microcapsules as an enteric delivery vehicle for probiotic bacteria. *Biomacromolecules*, 12 (7), 2834–2840.
- Dikit, P., H-Kittikun, A., Maneerat, S. (2015). Survival of encapsulated potentially probiotic *Lactobacillus plantarum* D6SM3 with bioemulsifier derived from spent yeast in simulated gastrointestinal conditions. *Songklanakarin Journal of Science and Technology*, 37(4), 425-432.
- FAO/WHO (2002). Joint FAO/WHO (Food and Agriculture Organization/World Health Organization) working group report on drafting guidelines for the evaluation of probiotics in food. London/Ontario, Canada: FAO/WHO.1–11.
- Hattingh, A. L., Viljoen, B.C. (2001). Growth and survival of a probiotic yeast in dairy products. *Food Research International*, 34(9), 791-796.
- Heydari, S., Mortazavian, A.M., Ehsani, M.R., Mohammadifar, M.A., Ezzatpanah, H., Sohrabvandi, S. (2012). Biochemical, microbiological and sensory characteristics

- of probiotic yogurt containing various prebiotic or fiber compounds. *Italian Journal of Food Science*, 23, 153–163.
- Homayouni, A., Azizi, A., Ehsani, M.R., Yarmand, M.S., Razavi, S.H. (2008). Effect of microencapsulation and resistant starch on the probiotic survival and sensory properties of synbiotic ice cream. *Food Chemistry*, 111, 50–55.
- Huq, T., Khan, A., Khan, R. A., Riedl, B., Lacroix, M. (2013). Encapsulation of probiotic bacteria in biopolymeric system. *Critical Reviews in Food Science and Nutrition*, 53(9), 909-916.
- Karaolis, C., Botsaris, G., Pantelides, I., Tsaltas, D. (2013). Potential application of *Saccharomyces boulardii* as a probiotic in goat's yoghurt: survival and organoleptic effects. *International Journal of Food Science and Technology*, 48, 1445–1452.
- Krasaekoopt, W., Bhandari, B., Deeth, H. (2006). Survival of probiotics encapsulated in chitosan-coated alginate beads in yogurt from UHT and conventionally treated milks during storage. *Lebensm.-Wiss. U.-Technol.*, 39, 177-183.
- Kurtzman, C.P., Fell, J.W., Boekhout, T. (2011). The yeasts, a Taxonomic Study .5th ed. Elsevier B.V., Volume 1. 537 p.
- Lee, K.Y., Mooney, D.J. (2012). Alginate: properties and biomedical applications. *Progress in Polymer Science*, 37(1), 106–126.
- Mohammadi, R., Mortazavian, A.M., Khosrokhavar, R., Da Cruz, A.G. (2011). Probiotic ice cream: viability of probiotic bacteria and sensory properties. *Ann Microbiol*, 61, 411–424.
- Niamah, A. K. (2017). Physicochemical and Microbial Characteristics of Yogurt with Added *Saccharomyces Boulardii*. *Current Research in Nutrition and Food Science Journal*, 5(3), 300-307.
- Niamah, A. K., Al-Sahlany, S. T. G., Al-Manhel, A. J. (2016). Gum arabic uses as prebiotic in yogurt production and study effects on physical, chemical properties and survivability of probiotic bacteria during cold storage. *World Applied Sciences Journal*, 34(9), 1190-1196.
- Nousia, F.G., Androulakis, P.I., Fletouris, D.J. (2011). Survival of *Lactobacillus acidophilus* LMGP-21381 in probiotic ice cream and its influence on sensory acceptability. *International Journal of Dairy Technology*, 64, 130–136.
- Pandiyan, C. (2010). Development of synbiotic ice cream enriched whey protein concentrate. (Doctoral dissertation). Tamil Nadu Veterinary and Animal Sciences University, Chennai, Tamil Nadu, India.
- Soukoulis, C., Fisk, I.D., Bohn, T. (2014). Ice cream as a vehicle for incorporating health-promoting ingredients: conceptualization and overview of quality and storage stability. *Comprehensive Reviews in Food Science and Food Safety*, 13, 627-655.
- Zamora-vega, R., Ez-soto, J.L., Martinez-Flores, H.C., Folres-Magallo, N.R., Mun˜oz-Ruiz, C.V., Lez, J.V., Ortega, T.S. (2012). Effect of incorporating prebiotics in coating materials for the microencapsulation of *Sacharomyces boulardii*. *International Journal of Food Sciences and Nutrition*, 63(8), 930–935.

Acknowledgment

The authors would like to thank to Biotechnology Lab. / Department of Food Science/ College of Agriculture/ University of Basrah.