

Separation Procedures of Biotechnology Products

Dr. Sarmad Ghazi
Biotechnology
Ph.D. Lecture

Stage # 1. Solid-Liquid Separation:

The first step in product recovery is the separation of whole cells (cell biomass) and other insoluble ingredients from the culture broth (Note: If the desired product is an intracellular metabolite, it must be released from the cells before subjecting to solid-liquid separation). Some authors use the term harvesting of microbial cells for the separation of cells from the culture medium. Several methods are in use for solid-liquid separation. These include flotation, flocculation, filtration and centrifugation.

Flotation الطفو :

When a gas is introduced into the liquid broth, it forms bubbles. The cells and other solid particles get adsorbed on gas bubbles. These bubbles rise to the foam layer which can be collected and removed. The presence of certain substances, referred to as collector substances, facilitates stable foam formation e.g., long chain fatty acids, amines.

Flocculation التلبد :

In flocculation, the cells (or cell debris) form large aggregates to settle down for easy removal. The process of flocculation depends on the nature of cells and the ionic constituents of the medium. Addition of flocculating agents (inorganic salt, organic polyelectrolyte, mineral hydrocolloid) is often necessary to achieve appropriate flocculation.

Filtration:

Filtration is the most commonly used technique for separating the biomass and culture filtrate. The efficiency of filtration depends on many factors— the size of the organism, presence of other organisms, viscosity of the medium, and temperature. Several filters such as depth filters, absolute filters, rotary drum vacuum filters and membrane filters are in use.

Membrane Filters:

In this type of filtration, membranes with specific pore sizes can be used. However, clogging of filters is a major limitation. There are two types of membrane filtrations—static filtration and cross-flow filtration (Fig. 20.3). In cross-flow filtration, the culture broth is pumped in a crosswise fashion across the membrane. This reduces the clogging process and hence better than the static filtration.




Fig. 20.3 : Filter systems for separation of biomass and culture filtrate (A) Static-flow filtration (B) Cross-flow filtration.

Types of filtration processes:

There are 3 major types of filtrations based on the particle sizes and other characters (Table 20.1). These are microfiltration, ultrafiltration and reverse osmosis.

TABLE 20.1 Major types of filtration processes with characteristic features

<i>Type</i>	<i>Sizes of particles separated</i>	<i>Compound or particle separated</i>
1. Microfiltration	0.1–10 μm	Cells or cell fractions, viruses.
2. Ultrafiltration	0.001–0.1 μm	Compounds with molecular weights greater than 1000 (e.g. enzymes).
3. Reverse osmosis (hyperfiltration)	0.0001–0.001 μm	Compounds with molecular weights less than 1000 (e.g. lactose).

Centrifugation:

The technique of centrifugation is based on the principle of density differences between the particles to be separated and the medium. Thus, centrifugation is mostly used for separating solid particles from liquid phase (fluid/particle separation). Unlike the centrifugation that is conveniently carried out in the laboratory scale, there are certain limitations for large scale industrial centrifugation.

However, in recent years, continuous flow industrial centrifuges have been developed. There is a continuous feeding of the slurry and collection of clarified fluid, while the solids deposited can be removed intermittently. The different types of centrifuges are depicted in Fig. 20.4, and briefly described hereunder.




Fig. 20.4 : Centrifuges commonly used in downstream processing (A) Tubular bowl centrifuge (B) Disc centrifuge (C) Multichamber centrifuge (D) Scroll centrifuge (decanter).

Stage # 2. Release of Intracellular Products:

As already stated, there are several biotechnological products (vitamins, enzymes) which are located within the cells. Such compounds have to be first released (maximally and in an active form) for their further processing and final isolation. The microorganisms or other cells can be disintegrated or disrupted by physical, chemical or enzymatic methods. The outline of different techniques used for breakage of cells is given in Fig. 20.5.




Fig. 20.5 : Major methods for cell disruption to release the intracellular products
(* indicate mechanical methods while all the remaining are non-mechanical).

The selection of a particular method depends on the nature of the cells, since there is a wide variation in the property of cell disruption or breakage. For instance, Gram-negative bacteria and filamentous fungi can be more easily broken compared to Gram-positive bacteria and yeasts.

Cell Disruption:

Physical methods of cell disruption:

The microorganisms or cells can be disrupted by certain physical methods to release the intracellular products.

Ultra sonication:

Ultrasonic disintegration is widely employed in the laboratory. However, due to high cost, it is not suitable for large-scale use in industries.

Osmotic shock:

This method involves the suspension of cells (free from growth medium) in 20% buffered sucrose. The cells are then transferred to water at about 4°C. Osmotic shock is used for the release of hydrolytic enzymes and binding proteins from Gram-negative bacteria.

Heat shock (thermolysis):

Breakage of cells by subjecting them to heat is relatively easy and cheap. But this technique can be used only for a very few heat-stable intracellular products.

High pressure homogenization:

This technique involves forcing of cell suspension at high pressure through a very narrow orifice to come out to atmospheric pressure. This sudden release of high pressure creates a liquid shear that can break the cells.

Impingement:

In this procedure, a stream of suspended cells at high velocity and pressure are forced to hit either a stationary surface or a second stream of suspended cells (impinge literally means to strike or hit). The cells are disrupted by the forces created at the point of contact. Micro fluidizer is a device developed based on the principle of impingement. It has been successfully used for breaking E. coli cells. The advantage with impingement technique is that it can be effectively used for disrupting cells even at a low concentration.

Grinding with glass beads:

The cells mixed with glass beads are subjected to a very high speed in a reaction vessel. The cells break as they are forced against the wall of the vessel by the beads. Several factors influence the cell breakage-size and quantity of the glass beads, concentration and age of cells, temperature and agitator speed. Under optimal conditions, one can expect a maximal breakage of about 80% of the cells.

A diagrammatic representation of a cell disrupter employing glass beads is shown in Fig. 20.6. It contains a cylindrical body with an inlet, outlet and a central motor-driven shaft. The cylinder is fitted with glass beads. The cell suspension is added through the inlet and the disrupted cells come out through the outlet. The body of the cell disrupter is kept cool while the operation is on.




Fig. 20.6 : Diagrammatic representation of a cell disrupter.

Mechanical and non-mechanical methods:

Among the physical methods of cell disruption described before, ultra sonication, high-pressure homogenization, impingement and grinding with glass beads are mechanical while osmotic shock and heat shock are non-mechanical. The chemical and enzymatic methods (described below) are non-mechanical in nature.

Chemical methods of cell disruption:

Treatment with alkalies, organic solvents and detergents can lyse the cells to release the contents.

Alkalies:

Alkali treatment has been used for the extraction of some bacterial proteins. However, the alkali stability of the desired product is very crucial for the success of this method e.g., recombinant growth hormone can be efficiently released from E. coli by treatment with sodium hydroxide at pH 11.

Organic solvents:

Several water miscible organic solvents can be used to disrupt the cells e.g., methanol, ethanol, isopropanol, butanol. These compounds are inflammable; hence require specialized equipment for fire safety. The organic solvent toluene is frequently used. It is believed that toluene dissolves membrane phospholipids and creates membrane pores for release of intracellular contents.

Detergents:

Detergents that are ionic in nature, cationic-cetyl trimethyl ammonium bromide or anionic-sodium lauryl sulfate can denature membrane proteins and lyse the cells. Non-ionic detergents (although less reactive than ionic ones) are also used to some extent e.g., Triton X-100 or Tween. The problem with the use of detergents is that they affect purification steps, particularly the salt precipitation. This limitation can be overcome by using ultrafiltration or ion-exchange chromatography for purification.

Enzymatic methods of cell disruption:

Cell disruption by enzymatic methods has certain advantages i.e., lysis of cells occurs under mild conditions in a selective manner. This is quite advantageous for product recovery. Lysozyme is the most frequently used enzyme and is commercially available (produced from hen egg white). It hydrolyses β -1, 4-glycosidic bonds of the mucopeptide in bacterial cell walls. The Gram- positive bacteria (with high content of cell wall mucopeptides) are more susceptible for the action of lysozyme.

For Gram-negative bacteria, lysozyme in association with EDTA can break the cells. As the cell wall gets digested by lysozyme, the osmotic effects break the periplasmic membrane to release the intracellular contents. Certain other enzymes are also used, although less frequently, for cell disruption. For the lysis of yeast cell walls, glucanase and mannanase in combination with proteases are used.

Combination of methods:

In order to increase the efficiency of cell disintegration in a cost-effective manner, a combination of physical, chemical and enzymatic methods is employed.

Stage # 3. Concentration:

The filtrate that is free from suspended particles (cells, cell debris etc.) usually contains 80-98% of water. The desired product is a very minor constituent. The water has to be removed to achieve the product concentration. The commonly used techniques for concentrating biological products are evaporation, liquid-liquid extraction, membrane filtration, precipitation and adsorption. The actual procedure adopted depends on the nature of the desired product (quality and quantity to be retained as far as possible) and the cost factor.

Evaporation:

Water in the broth filtrate can be removed by a simple evaporation process. The evaporators, in general, have a heating device for supply of steam, and unit for the separation of concentrated product and vapour, a condenser for condensing vapour, accessories and control equipment. The capacity of the equipment is variable that may range from small laboratory scale to industrial scale.

Liquid-Liquid Extraction:

The concentration of biological products can be achieved by transferring the desired product (solute) from one liquid phase to another liquid phase, a phenomenon referred to as liquid-liquid extraction. Besides concentration, this technique is also useful for partial purification of a product.

The efficiency of extraction is dependent on the partition coefficient i.e. the relative distribution of a substance between the two liquid phases. The process of liquid-liquid extraction may be broadly categorized as extraction of low molecular weight products and extraction of high molecular weight products.

Extraction of low molecular weight products:

By using organic solvents, the lipophilic compounds can be conveniently extracted. However, it is quite difficult to extract hydrophilic compounds. Extraction of lipophilic products can be done by the following techniques.

Physical extraction:

The compound gets itself distributed between two liquid phases based on the physical properties. This technique is used for extraction of non-ionising compounds.

Dissociation extraction:

This technique is suitable for the extraction of ionisable compounds. Certain antibiotics can be extracted by this procedure.

Reactive extraction:

In this case, the desired product is made to react with a carrier molecule (e.g., phosphorus compound, aliphatic amine) and extracted into organic solvent. Reactive extraction procedure is quite useful for the extraction of certain compounds that are highly soluble in water (aqueous phase) e.g., organic acids.

Membrane Filtration:

Membrane filtration has become a common separation technique in industrial biotechnology. It can be conveniently used for the separation of biomolecules and particles, and for the concentration of fluids. The membrane filtration technique basically involves the use of a semipermeable membrane that selectively retains the particles/molecules that are bigger than the pore size while the smaller molecules pass through the membrane pores.

Membranes used in filtration are made up of polymeric materials such as polyethersulfone and polyvinyl difluoride. It is rather difficult to sterilize membrane filters. In recent years, micro-filters and ultrafilters composed of ceramics and steel are available. Cleaning and sterilization of such filters are easy.

Membrane adsorbers:

They are micro- or macro porous membranes with ion exchange groups and/or affinity ligands. Membrane adsorbers can bind to proteins and retain them. Such proteins can be eluted by employing solutions in chromatography.

Precipitation:

Precipitation is the most commonly used technique in industry for the concentration of macromolecules such as proteins and polysaccharides. Further, precipitation technique can also be employed for the removal of certain unwanted byproducts e.g. nucleic acids, pigments.

Neutral salts, organic solvents, high molecular weight polymers (ionic or non-ionic), besides alteration in temperature and pH are used in precipitation. In addition to these non-specific protein precipitation reactions (i.e. the nature of the protein is unimportant), there are some protein specific precipitations e.g., affinity precipitation, ligand precipitation.

Neutral salts:

The most commonly used salt is ammonium sulfate, since it is highly soluble, nontoxic to proteins and low-priced. Ammonium sulfate increases hydrophobic interactions between protein molecules that result in their precipitation. The precipitation of proteins is dependent on several factors such as protein concentration, pH and temperature.

Organic solvents:

Ethanol, acetone and propanol are the commonly used organic solvents for protein precipitation. They reduce the dielectric constant of the medium and enhance electrostatic interaction between protein molecules that lead to precipitation. Since proteins are denatured by organic solvents, the precipitation process has to be carried out below 0° C.

Increase in temperature:

The heat sensitive proteins can be precipitated by increasing the temperature.

Change in pH:

Alterations in pH can also lead to protein precipitation.

Affinity precipitation:

The affinity interaction (e.g., between antigen and antibody) is exploited for precipitation of proteins.

Adsorption:

The biological products of fermentation can be concentrated by using solid adsorbent particles. In the early days, activated charcoal was used as the adsorbent material. In recent years, cellulose-based adsorbents are employed for protein concentration.

And for concentration of low molecular weight compounds (vitamins, antibiotics, peptides) polystyrene, methacrylate and acrylate based matrices are used. The process of adsorption can be carried out by making a bed of adsorbent column and passing the culture broth through it. The desired product, held by the adsorbent, can be eluted.

Stage # 4. Purification by Chromatography:

The biological products of fermentation (proteins, pharmaceuticals, diagnostic compounds and research materials) are very effectively purified by chromatography. It is basically an analytical technique dealing with the separation of closely related compounds from a mixture. Chromatography usually consists of a stationary phase and mobile phase.

The stationary phase is the porous solid matrix packed in a column (equilibrated with a suitable solvent) on to which the mixture of compounds to be separated is loaded. The compounds are eluted by a mobile phase.

A single mobile phase may be used continuously or it may be changed appropriately to facilitate the release of desired compounds. The eluate from the column can be monitored continuously (e.g. protein elution can be monitored by ultraviolet adsorption at 280 nm), and collected in fractions of definite volumes.

The different types of chromatography techniques used for separation (mainly proteins) along with the principles are given in Table 20.2. A large number of matrices are commercially available for purification of proteins e.g., agarose, cellulose, polyacrylamide, porous silica, cross-linked dextran, polystyrene. Some of the important features of selected chromatographic techniques are briefly described.

TABLE 20.2 Chromatographic techniques along with the principles for separation of proteins

<i>Chromatography</i>	<i>Principle</i>
Gel-filtration (size exclusion)	Size and shape
Ion-exchange	Net charge
Chromatofocussing	Net charge
Affinity	Biological affinity and molecular recognition
Hydrophobic interaction	Polarity (hydrophobicity of molecules)
Immobilized metal-ion affinity	Metal ion binding

Gel-filtration chromatography:

This is also referred to as size-exclusion chromatography. In this technique, the separation of molecules is based on the size, shape and molecular weight. The sponge-like gel beads with pores serve as molecular sieves for separation of smaller and bigger molecules. A solution mixture containing molecules of different sizes (e.g. different proteins) is applied to the column and eluted.

The smaller molecules enter the gel beads through their pores and get trapped. On the other hand, the larger molecules cannot pass through the pores and therefore come out first with the mobile liquid (Fig. 20.7). At the industrial scale, gel-filtration is particularly useful to remove salts and low molecular weight compounds from high molecular weight products.




Fig. 20.7 : The principle of gel-filtration chromatography.

Ion-exchange chromatography:

It involves the separation of molecules based on their surface charges. Ion-exchangers are of two types (cation-exchangers which have negatively charged groups like carboxymethyl and sulfonate, and anion-exchangers with positively charged groups like diethylaminoethyl (DEAE)).

In ion-exchange chromatography, the pH of the medium is very crucial, since the net charge varies with pH. In other words, the pH determines the effective charge on both the target molecule and the ion-exchanger. The ionic bound molecules can be eluted from the matrix by changing the pH of the eluant or by increasing the concentration of salt solution. Ion-exchange chromatography is useful for the purification of antibiotics, besides the purification of proteins.

Affinity chromatography:

This is an elegant method for the purification of proteins from a complex mixture. Affinity chromatography is based on an interaction of a protein with an immobilized ligand. The ligand can be a specific antibody, substrate or an inhibitor. The immobilized ligand on a solid matrix can be effectively used to fish out complementary structures.

In Table 20.3, some examples of ligands used for the purification of proteins are given. The protein bound to the ligand can be eluted by reducing their interaction. This can be achieved by changing the pH of the buffer, altering the ionic strength or by using another free ligand molecule. The fresh ligand used has to be removed in the subsequent steps.

TABLE 20.3 Some examples of ligands used for separation of proteins by affinity chromatography

<i>Ligand</i>	<i>Type of protein</i>
Antibody	Antigen
Cofactor	Enzyme
Receptor	Hormone
Hapten	Antibody
Inhibitor	Enzyme
Lectins	Glycoproteins
Heparin	Coagulation factors
Metal ions	Metal ion binding proteins

Stage # 5. Formulation:

Formulation broadly refers to the maintenance of activity and stability of a biotechnological products during storage and distribution. The formulation of low molecular weight products (solvents, organic acids) can be achieved by concentrating them with removal of most of the water. For certain small molecules, (antibiotics, citric acid), formulation can be done by crystallization by adding salts.

Proteins are highly susceptible for loss of biological activity; hence their formulation requires special care. Certain stabilizing additives are added to prolong the shelf life of protein. The stabilizers of protein formulation include sugars (sucrose, lactose), salts (sodium chloride, ammonium sulfate), polymers (polyethylene glycol) and polyhydric alcohols (glycerol). Proteins may be formulated in the form of solutions, suspensions or dry powders.

Drying:

Drying is an essential component of product formulation. It basically involves the transfer of heat to a wet product for removal of moisture. Most of the biological products of fermentation are sensitive to heat, and therefore require gentle drying methods. Based on the method of heat transfer, drying devices may be categorized as contact, convection, radiation dryers. These three types of dryers are commercially available.

Spray drying:


Spray drying is used for drying large volumes of liquids. In spray drying, small droplets of liquid containing the product are passed through a nozzle directing it over a stream of hot gas. The water evaporates and the solid particles are left behind.

Freeze-drying:


Freeze-drying or lyophilization is the most preferred method for drying and formulation of a wide-range of products—pharmaceuticals, foodstuffs, bacteria, viruses. This is mainly because freeze-drying usually does not cause loss of biological activity of the desired product.

Lyophilization is based on the principle of sublimation of a liquid from a frozen state. In the actual technique, the liquid containing the product is frozen and then dried in a freeze-dryer under vacuum. The vacuum can now be released and the product containing vials can be sealed e.g., penicillin can be freeze dried directly in ampules.

Gel Filtration



Ion Exchange



Affinity

