Chapter one

Functions

Sets of Numbers:

- $\mathbb{N} = \{\mathbf{0}, \mathbf{1}, \mathbf{2}, \dots\}$ Natural Numbers.
- $\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ Integer Numbers.
- $\mathbb{Q} = \{\frac{a}{b} : a, b \in \mathbb{Z}, b \neq 0\}$ Rational Numbers.
- $\mathbb{IQ} = \pi, e, \sqrt{2}, \sqrt{3}, \sqrt{5}, \dots$ Irrational Numbers.

 $\mathbb{R} = \mathbb{Q} \cup \mathbb{I}\mathbb{Q} \qquad \text{Real Numbers}$

Definition: If *a* and *b* are real numbers, we define the Intervals as follows:

- (1) Open Intervals $(a,b) = \{x \in \mathbb{R} : a < x < b\}$
- (2) Closed Intervals $[a,b] = \{x \in \mathbb{R} : a \le x \le b\}$
- (3) Half open, half closed Intervals $[a,b] = \{x \in R : a \le x < b\}$ and $(a,b] = \{x \in R : a < x \le b\}.$
- (4) $[a,\infty) = \{x \in R : x \ge a\}, (-\infty, a] = \{x \in R : x \le a\}, (-\infty, a) = \{x \in R : x < a\}$ and $(-\infty, \infty) = \{x \in R : x \text{ is } a \text{ real number}\} = R.$

<u>Remark</u>: If *A* and *B* are intervals, then

- The union of A and B is denoted by A ∪ B and defined as the interval whose members belong to A or B (or both).
- 2. The intersection of *A* and *B* is denoted by $A \cap B$ and defined as the interval whose members belong to both *A* and *B*.

Example: Let A = [0,5] and B = [1,7] then $A \cup B = [0,7]$ and $A \cap B = [1,5]$.

Functions

<u>Definition</u>: A relation $f: X \to Y$ is called a function if and only if for each element $x \in X$ there exist a unique element $y \in Y$ such that y = f(x).

Note:

(3)

(4)

- (1) x is the independent variable (input value of f) and y is dependent variable (output value of f at x)
- (2) The set X of all possible input values is called the domain of f and it's denoted by D_f
- (3) The set *Y* is called the co-domain of the function.
- (4) The set of all possible output values f(x) as x varies throughout D_x is called the range of f and it's denoted by R_f . (Note that $R_f \subseteq Y$.)

Examples: Find the Domain and the Range of the following:

(1)
$$y = x + 5$$

$$D_f = \{x \in R : -\infty < x < \infty\} = R$$

To find the Range, we represent x in terms of y. x = y - 5 $R_f = \{y \in R : -\infty < y < \infty\} = R$

(2)
$$y = x^2$$
 $D_f = \{x \in R : -\infty < x < \infty\} = R$

$$\boldsymbol{x} = \sqrt{\boldsymbol{y}} \implies R_f = \{ \boldsymbol{y} \in \boldsymbol{R} : \boldsymbol{y} \ge 0 \} = [0, \infty)$$

(set the denominator = 0)

 $y = \frac{1}{x+2}$ $x+2 = 0, \quad x = -2$

$$x = \frac{1-2y}{y} \implies D_f = \{x \in \Re : x \neq -2\} = \Re/\{-2\}$$
$$\Rightarrow R_f = \{y \in \Re : y \neq 0\} = \Re/\{0\}$$
$$y = \sqrt{x+9} \qquad x \in [0,7] \qquad \Rightarrow D_f = [0,7]$$

Put x = 0 in the function we get y = 3Put x = 7 in the function we get y = 4 then $R_f = [3,4]$

(5)
$$f(x) = \frac{1}{\sqrt{2-x}} + 5$$
 (6) $y = \frac{3x}{x^2 - 5x + 6}$

Algebraic Combination of Function

If f and g are two functions with domains D_f and D_g respectively, then

- (1) (f+g)(x) = f(x) + g(x) with $D_{f+g} = D_f \cap D_g$
- (2) (f-g)(x) = f(x) g(x) with $D_{f-g} = D_f \cap D_g$
- (3) $(f.g)(x) = f(x) \cdot g(x)$ with $D_{f.g} = D_f \cap D_g$
- (4) $((\frac{f}{g})(x) = \frac{f(x)}{g(x)}, \quad g(x) \neq 0 \text{ with } D_{f/g} = D_f \cap D_g \text{ and } g(x) \neq 0$

Examples: 1) If $f(x) = x^2 - 5x - 6$ and $g(x) = 3x^2 + 4$ then find (f + g)(x), $(f - g)(x), (g - f)(x), (f \cdot g)(x), (f/g)(x)$ and (g/f)(x). Moreover, find their domains.

Solution: $D_f = \mathbb{R}$, $D_a = \mathbb{R}$

Function	Formula	Domain
f + g	$(f+g)(x) = 4x^2 - 5x - 2$	R
f-g	$(f-g)(x) = -2x^2 - 5x - 10$	R
g-f	$(g-f)(x) = 2x^2 + 5x + 10$	R
f.g	$(f.g)(x) = 3x^4 - 15x^3 - 14x^2 - 20x$	R
	- 24	
f/g	$\left(\frac{f}{g}\right)(x) = \frac{x^2 - 5x - 6}{3x^2 + 4}$	R
g/f	$\left(\frac{g}{f}\right)(x) = \frac{3x^2 + 4}{x^2 - 5x - 6}$	$(-\infty, -1) \cup (-1, 6) \cup (6, \infty)$

2) If $f(x) = \sqrt{x}$ and $gx) = \sqrt{1-x}$ then find (f+g)(x), (f-g)(x), (g-f)(x), (f,g)(x), (f/g)(x) and (g/f)(x). Moreover, find their domains.

Solution: D _f	$= \{x \in \mathbb{R} : x \ge 0\} = \mathbb{R}^+ \cup \{0\} = [0, \infty), D_g = [0, \infty]$	$= \{x \in \mathbb{R} : x \le 1\} = (-\infty, 1]$
Function	Formula	Domain
f + g	$(f+g)(x) = \sqrt{x} + \sqrt{1-x}$	[0,1]
f-g	$(f-g)(x) = \sqrt{x} - \sqrt{1-x}$	[0,1]
g-f	$(g-f)(x) = \sqrt{1-x} - \sqrt{x}$	[0,1]
<i>f</i> . <i>g</i>	$(f.g)(x) = \sqrt{x(1-x)}$	[0,1]
f/g	$\left(\frac{f}{g}\right)(x) = \frac{\sqrt{x}}{\sqrt{1-x}} = \sqrt{\frac{x}{1-x}}$	[0,1)
g/f	$\left(\frac{g}{f}\right)(x) = \frac{\sqrt{1-x}}{\sqrt{x}} = \sqrt{\frac{1-x}{x}}$	(0,1]

Department of Mathematics - College of Education for Pure Sciences - University of Basrah Telegram: @uobmath t.me\uobmath 3

3) If
$$f(x) = 2$$
 and $gx) = x^2$ then find $(f + g)(x)$, $(f - g)(x)$, $(g - f)(x)$, $(f \cdot g)(x)$, $(f/g)(x)$ and $(g/f)(x)$. Moreover, find their domains and ranges.

Function	Formula	Domain	Rang
f + g	$(f+g)(x) = 2 + x^2$	R	[2,∞)
f-g	$(f-g)(x) = 2 - x^2$	\mathbb{R}	(−∞,2)
g-f	$(g-f)(x) = x^2 - 2$	\mathbb{R}	[−2,∞)
<i>f</i> . <i>g</i>	$(f.g)(x) = 2x^2$	\mathbb{R}	[0,∞)
f/g	$\left(\frac{f}{g}\right)(x) = \frac{2}{x^2}$	$(-\infty,0) \cup (0,\infty)$	(0,∞)
g/f	$\left(\frac{g}{f}\right)(x) = \frac{x^2}{2}$	R	(0,∞)

Solution: $D_f = \mathbb{R}$, $D_g = \mathbb{R}$

Composite Function

If the range of the function g(x) is contained in the domain of the function f(x), then the composition $f \circ g$ is the function defined by $(f \circ g)(x) = f(g(x))$. The domain of $f \circ g$ consists of the number x in the domain of g for which g(x) lies in the domain of f.

$$x \xrightarrow{g} g(x) \xrightarrow{f} f(g(x))$$

 $f \circ g$

g(x)

Examples:

1) If $f(x) = \sqrt{x}$ and g(x) = x + 1, then find (a) $(f \circ g)(x)$ (b) $(g \circ f)(x)$ (c) $(f \circ f)(x)$ (d) $(g \circ g)(x)$

Solution: $D_f = [0, \infty), D_g = \mathbb{R}$ CompositeDomain(a) $(f \circ g)(x) = f(g(x)) = \sqrt{g(x)} = \sqrt{x+1}$ $[-1, \infty)$ (b) $(g \circ f)(x) = g(f(x)) = f(x) + 1 = \sqrt{x} + 1$ $[0, \infty)$ (c) $(f \circ f)(x) = f(f(x)) = \sqrt{f(x)} = \sqrt{\sqrt{x}} = x^{\frac{1}{4}}$ $[0, \infty)$ (d) $(g \circ g)(x) = g(g(x)) = g(x) + 1 = (x+1) + 1 = x + 2$ \mathbb{R}

2) If $f(x) = x^2 - 3$ and g(x) = x + 1, then find (a) $(f \circ g)(x)$ (b) $(g \circ f)(x)$ (c) $(f \circ f)(x)$ (d) $(g \circ g)(x)$ Solution: $D_f = \mathbb{R}$, $D_g = \mathbb{R}$

Composite	Domain
(a) $(f \circ g)(x) = f(g(x)) = (g(x))^2 - 3 = x^2 + 2x - 2$	\mathbb{R}
(b) $(g \circ f)(x) = g(f(x)) = f(x) + 1 = x^2 - 2$	\mathbb{R}
$(c) (f \circ f)(x) = f(f(x)) = (f(x))^2 - 3 = (x^2 - 3)^2 - 3 = x^4 - 6x^2 + 6$	R
(d) $(g \circ g)(x) = g(g(x)) = g(x) + 1 = (x + 1) + 1 = x + 2$	\mathbb{R}

The Inverse of the functions:

<u>Definition</u>: A function is said to be one - to - one function if and only if there is no two elements of the domain have the same image in the range.



Horizontal Line Test: A function is one-to-one if and only if there is no horizontal line can intersect its graph more than once.



Definition: Let f be a one-to-one function with domain A and range B (bijective function), then its inverse function f^{-1} has domain B and range A and is defined by $f^{-1}(y) = x \Leftrightarrow f(x) = y, \forall y \in B$ or $f^{-1}(f(x)) = f(f^{-1}(x)) = x$ and $D_{f^{-1}} = R_f$, $D_f = R_{f^{-1}}$.

How to find the inverse function of a one-to-one function f

Step 1: Write y = f(x).

Step 2: Solve this equation for x in terms of y (if possible). **Step 3:** To express f^{-1} as a function of x, interchange x and y.

The resulting equation is $y = f^{-1}(x)$.

Example: Find the inverse of f for $f(x) = x^3 + 2$ **Solution:** According to the above algorithm, we first write $y = x^3 + 2$, then we solve this equation for x:

 $x^3 = y - 2 \Longrightarrow x = \sqrt[3]{y - 2}$ Finally, we iterchange x and y: $y = \sqrt[3]{x - 2}$

 $y = \sqrt{x - 2}$ $\therefore f^{-1}(x) = \sqrt[3]{x - 2}$

<u>CAUTION</u> The -1 in f^{-1} is not a power number. Thus $f^{-1}(x)$ dose not mean $\frac{1}{f(x)}$

✤ The graph of f^{-1} is obtained by reflecting the graph of f around the line y = x.



f(x)

Graph of Functions:

<u>Definition</u>: Let f(x) be a function with domain D_f . The set of all points (x, y) in the plane with x in D_f and y = f(x) is called the graph of f(x). { $(x, f(x)): x \in D_f$ }

x	1	2		n		f(1)	ſ	
f(x)	f(1)	f(2)		f(n)			f(2)	
(x,f(x))	(1, f(1))	(2, f(2))	•••	(n, f(n))				x
					0	1	2	 +
								÷.

<u>Shifting a Graph of a function:</u> 1. <u>Vertical Shifts</u>:

 $y = f(x) + k \Longrightarrow \text{Shifts the graph of } \begin{cases} up \ k \text{ units if } k > 0 \\ down \ |k| \text{ units if } k < 0 \end{cases}$

2. Horizontal Shifts:

 $y = f(x + h) \Rightarrow$ Shifts the graph of $f \begin{cases} left h units if h > 0 \\ right |h| units if h < 0 \end{cases}$



Scaling and Reflecting a Graph of a Function:

For c > 1, the graph is scaled as:

y = cf(x) Stretches the graph of *f* vertically by a factor of *c*. $y = \frac{1}{c}f(x)$ Compresses the graph of *f* vertically by a factor of *c*. y = f(cx) Compresses the graph of *f* horizontally by a factor of *c*. $y = f(\frac{1}{c}x)$ Stretches the graph of *f* horizontally by a factor of *c*.

For c = -1, the graph is reflected as:

y = -f(x)	Reflects the graph f across the $x - axis$
y = f(-x)	Reflects the graph f across the $y - axis$



<u>Some Specific Types of Functions</u> Algebraic Functions:

1. Real valued function:

The function $f: X \to Y$ is called a real valued function if $X \subseteq \mathbb{R}$ and $Y \subseteq \mathbb{R}$.

2. <u>Constant function</u>:

The function $f: \mathbb{R} \to \mathbb{R}$, which is defined by y = f(x) = c, $\forall x \in \mathbb{R}$, where *c* is constant in \mathbb{R} , is called a constant function. $D_f = \mathbb{R}$ and $R_f = \{c\}$. **Example:** Consider the function y = f(x) = 2

x	-2	-1	0	1	2
f(x)	2	2	2	2	2
$(x, \overline{f}(x))$	(-2,2)	(-1,2)	(0,2)	(1,2)	(2,2)

3. <u>Identity function</u>:

The function $f: \mathbb{R} \to \mathbb{R}$, which is defined by y = f(x) = x, $\forall x \in \mathbb{R}$, is called the identity function. $D_f = \mathbb{R}$ and $R_f = \mathbb{R}$.

x	-2	-1	0	1	2
f(x)	-2	-1	0	1	2
(x, f(x))	(-2, -2)	(-1, -1)	(0,0)	(1,1)	(2,2)

4. Power function:

The function $f: X \to Y$, which is defined by $y = f(x) = x^a$, $\forall x \in X$, is called the power function.

a) If $a = n \in \mathbb{N}$: graph for a = 1,2,3,4,5 $a = 1 \rightarrow f(x) = x$ (Identity function) $a = 2 \rightarrow f(x) = x^2 \rightarrow D_f = \mathbb{R}$ and $R_f = [0, \infty)$. $a = 3 \rightarrow f(x) = x^3 \rightarrow D_f = \mathbb{R}$ and $R_f = \mathbb{R}$. $a = 4 \rightarrow f(x) = x^4 \rightarrow D_f = \mathbb{R}$ and $R_f = [0, \infty)$. $a = 5 \rightarrow f(x) = x^5 \rightarrow D_f = \mathbb{R}$ and $R_f = \mathbb{R}$.





5. Polynomial function:

A polynomial function is any function can be written in the form

 $f(x) = P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0, n \in \mathbb{N}$ with the coefficients $a_n, a_{n-1}, \dots, a_2, a_1, a_0 \in \mathbb{R}$ and $D_f = \mathbb{R}$. with $a_n \neq 0, n \ge 0, (n \text{ is the degree of polynomial})$

- if $n=0 \rightarrow f(x) = P_0(x) = a_0$ (constant function)
- if $n=1 \rightarrow f(x) = P_1(x) = a_1 x + a_0$ (linear function)
- if $n \ge 2 \to f(x) = P_n(x) = a_n x^n + a_{n-1} x^{n-1} + \dots + a_2 x^2 + a_1 x + a_0$ (nonlinear function)

10

6. Absolute value function:

The function $f: \mathbb{R} \to \mathbb{R}$, which is defined by $y = f(x) = \begin{cases} x, & x \ge 0 \\ -x, & x < 0 \end{cases}$ is called the absolute value function (the modulus function). $D_f = \mathbb{R}$ and $R_f = \mathbb{R}^+ \cup \{0\} = [0, \infty)$.



7. Sign function

The function $f: \mathbb{R} \to \mathbb{R}$, which is defined by

$$f(x) = sgn(x) = \begin{cases} \frac{|x|}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases} = \begin{cases} 1 & \text{if } x > 0 \\ 0 & \text{if } x = 0 \\ -1 & \text{if } x < 0 \end{cases}$$

is called the sign function.

 $D_f = \mathbb{R}$ and $R_f = \{-1, 0, 1\}$



8. The Greatest integer function)

The function $f : \mathbb{R} \to \mathbb{R}$, which defined by

$$f(x) = [x] = \begin{cases} -2 & if -2 \le x < -1 \\ -1 & if -1 \le x < 0 \\ 0 & if \ 0 \le x < 1 \\ 1 & if \ 1 \le x < 2 \\ 2 & if \ 2 \le x < 3 \end{cases}$$

is called the greatest integer function. $D_f = \mathbb{R}$ and $R_f = \mathbb{Z}$.



Remark: Another name of the gratest integer function is the **Floor Function**.

Remark: There is another function called **Celling Function** denoted by [x] that maps x to the least integer, greater than or equal to x.

9. <u>The piecewise function</u>

A piecewise function is a function that is described by using different formulas on different part of its domain.



10. <u>Odd function</u>: The function $f: X \to Y$ is an odd function if f(-x) = -f(x).

11. <u>Even function</u>: The function $f: X \to Y$ is an even function if f(-x) = f(x).

12. <u>Rational function</u>: A rational function is a quotient or ratio $f(x) = \frac{p(x)}{q(x)}$, where p and q are polynomials. The domain of a rational function is the set of all real x for which $q(x) \neq 0$. The graphs of several rational functions are shown here below.



Transcendental Functions:

1. Trigonometric Functions:

Definition: A function f(x) is a periodic function if there is a positive number p such that f(x + p) = f(x) for every value of x. The positive number p is called the period of the function f. $sin(x + 2\pi) = sin(x), cos(x + 2\pi) = cos(x)$ $sec(x + 2\pi) = sec(x), csc(x + 2\pi) = csc(x)$ $period 2\pi$. $tan(x + \pi) = tan(x)$ $cot(x + \pi) = cot(x)$ $period \pi$.



Domain: $-\infty < x < \infty$ Range: $-1 \le y \le 1$ Period: 2π



Domain: $x \neq \pm \frac{\pi}{2}, \pm \frac{3\pi}{2}, \dots$ Range: $-\infty < y < \infty$ Period: π





Domain: $-\infty < x < \infty$ Range: $-1 \le y \le 1$ Period: 2π



Domain: $x \neq 0, \pm \pi, \pm 2\pi, \dots$ Range: $-\infty < y < \infty$ Period: π



University of Basrah College of Education for Pure Sciences Department of Mathematics

Some Important Identities of The Trigonometric Functions

 $\begin{aligned} \tan x &= \frac{\sin x}{\cos x} \\ \cot x &= \frac{\cos x}{\sin x} \\ \sec x &= \frac{1}{\cos x} \\ \sec x &= \frac{1}{\sin x} \\ \sin^2 x + \cos^2 x &= 1 \\ \sin 2x &= 2\sin x \cos x \\ \cos 2x &= \cos x \cos x - \sin x \sin x = \cos^2 x - \sin^2 x = 2\cos^2 x - 1 = 1 - 2\sin^2 x \\ \tan 2x &= \frac{2\tan x}{1 - \tan^2 x} \\ \sin(x \pm y) &= \sin x \cos y \pm \cos x \sin y \\ \cos(x \pm y) &= \cos x \cos y \mp \sin x \sin y \\ \tan(x \pm y) &= \frac{\tan x \pm \tan y}{1 \mp \tan x \tan y} \end{aligned}$

Definition by Power series

$$\sin x = x - rac{x^3}{3!} + rac{x^5}{5!} - rac{x^7}{7!} + \dots = \sum_{n=0}^\infty rac{(-1)^n x^{2n+1}}{(2n+1)!} \ \cos x = 1 - rac{x^2}{2!} + rac{x^4}{4!} - rac{x^6}{6!} + \dots = \sum_{n=0}^\infty rac{(-1)^n x^{2n}}{(2n)!}$$

Remark: One can define tangent, cotangent, secant and cosecant using identities above.

Inverse of Trigonometric Functions

The six basic trigonometric functions are not one-to-one (their values repeat periodically). However, we can restrict their domains to intervals on which they are one-to-one. The sine function increases from -1 at $x = -\pi/2$ to +1 at $x = \pi/2$. By restricting its domain to the interval $[-\pi/2, \pi/2]$, we make it one-to-one, so that it has an inverse $\sin^{-1} x$. Similar domain restrictions can be applied to all six trigonometric functions.

Domain restrictions that make the trigonometric functions one-to-one



University of Basrah College of Education for Pure Sciences Department of Mathematics



Graphs of the six basic inverse trigonometric functions.

2. Exponential Functions:

The function of the form $f(x) = a^x$ where a > 0 and $a \neq 1$, is called an exponential function. The domain of the exponential function is \mathbb{R} .



University of Basrah College of Education for Pure Sciences Department of Mathematics

Examples: 1) $y = f(x) = 2^x \rightarrow D_f = \mathbb{R}, R_f = (0, \infty)$

х	$f(x) = 2^x$
-2	$\mathbf{f}(-2) = 2^{-2} = \frac{1}{2^2} = \frac{1}{4}$
-1	$\mathbf{f}(-1) = 2^{-1} = \frac{1}{2^1} = \frac{1}{2}$
0	$f(0) = 2^0 = 1$
1	$f(1) = 2^1 = 2$
2	$f(2) = 2^2 = 4$



2)
$$y = f(x) = \left(\frac{1}{2}\right)^{x} \rightarrow D_{f} = \mathbb{R}, R_{f} = (0, \infty)$$

$$\begin{array}{c|c} x & f(x) = \left(\frac{1}{2}\right)^{x} \\ \hline -2 & f(-2) = \left(\frac{1}{2}\right)^{-2} = \frac{1^{-2}}{2^{-2}} = \frac{2^{2}}{1^{2}} = 4 \\ \hline -1 & f(-1) = \left(\frac{1}{2}\right)^{-1} = \frac{1^{-1}}{2^{-1}} = \frac{2^{1}}{1^{1}} = 2 \\ \hline 0 & f(0) = \left(\frac{1}{2}\right)^{0} = 1 \\ \hline 1 & f(1) = \left(\frac{1}{2}\right)^{1} = \frac{1^{1}}{2^{1}} = \frac{1}{2} \\ \hline 2 & f(2) = \left(\frac{1}{2}\right)^{2} = \frac{1^{2}}{2^{2}} = \frac{1}{4} \end{array}$$

17



3. Logarithmic Functions:

The function of the form $f(x) = log_a(x)$ where a > 0 and $a \neq 1$, called **logarithmic** function. The domain of **logarithmic function** is \mathbb{R}^+ . $y = log_a(x) \Leftrightarrow a^y = x$

Lows of logarithms:

If x and y are real numbers, then 1. $log_a(x, y) = log_a(x) + log_a(y)$ 2. $log_a(x/y) = log_a(x) - log_a(y)$ 3. $log_a(x^r) = r \cdot log_a(x)$ (where $r \in \mathbb{R}$)



The Natural Logarithmic Function:

The function of the form $f(x) = log_e(x) = ln(x)$, x > 0 called the natural logarithmic function. $D_f = \mathbb{R}^+$, $R_f = \mathbb{R}$.



<u>Exercises 2</u> In exercises (1-20), find the domain and range of each the following where					
y = f(x)					
1. $y = 5x + 3$	2. $y = 2x^2 + 1$	3. $y = -7x - 4$			
4. <i>y</i> = 7	5. $y = 4 - x^2$	6. $y = \sqrt{5x + 10}$			
7. $y = \sqrt{x^2 - 3x}$	8. $y = 1 - \sqrt{x}$	9. $y = \sqrt{x+9}$			
10. $y = \sqrt{3x - 4}$	11. $y = \sqrt{x^2 - 4}$	12. $y = \sqrt{4 - x^2}$			
13. $y = \frac{1}{7-x}$	14. $y = \frac{6}{x+2}$	15. $y = \sqrt{\frac{1}{x-2}}$			
16. $y = \frac{2}{\sqrt{2x-5}}$	17. $y = \frac{x}{\sqrt{x+1}}$	18. $y = \frac{1}{1 - \frac{1}{x - 2}}$			
19. $y = 2 + \frac{x^2}{x^2 + 4}$	20. $y = \frac{2}{x^2 - 16}$				

In exercises (21-35), find the domain and the range and sketch the graph of each the following function

 $21. y = \sqrt{x-5} \qquad 22. y = |2x+1| \qquad 23. y = \frac{3x+|x|}{x}$ $24. y = \frac{|x|}{x^2} \qquad 25. y = \begin{cases} x & if \ x \le 0 \\ x+1 & if \ x > 0 \end{cases} \qquad 26. y = \begin{cases} 2x+3 & if \ x < -1 \\ 3-x & if \ x \ge -1 \end{cases}$ $27. y = \begin{cases} x+2 & if \ x \le -1 \\ x^2 & if \ x > -1 \end{cases} \qquad 28. \begin{cases} -1 & if \ x \le -1 \\ 3x+2 & if \ |x| < 1 \\ 7-2x & if \ x \ge 1 \end{cases} \qquad 29. y = \begin{cases} \frac{1}{x} & if \ x < 0 \\ x & if \ x \ge 0 \end{cases}$ $30. y = \frac{1}{|x|} \qquad 31. y = |x^2 - 1| \qquad 32. y = [\sin(x)]$

33. $y = [\cos(x)]$ 34. $y = \sqrt{|x|}$ 35. $y = \frac{x^2}{|x|}$

In exercises 36 and 37, write formulas for $f \circ g$ and $g \circ f$ and find the domain and the range of each

36. $f(x) = \sqrt{x+1}$, $g(x) = \frac{1}{x}$ 37. $f(x) = x^2$, $g(x) = 1 - \sqrt{x}$

- 38. Let $f(x) = \frac{x}{x-2}$. Find a function y = g(x) so that $(f \circ g)(x) = x$.
- 39. Let $f(x) = 2x^3 4$. Find a function y = g(x) so that $(f \circ g)(x) = x + 2$.

In exercises 40-54, graph each function, not by plotting points, but by applying appropriate transformation to the graph of the standard functions.

$40. \ y = -\sqrt{2x+1}$	$41. \ y = \sqrt{1 - \frac{x}{2}}$	42. $y = (x - 1)^3 + 2$
43. $y = (1 - x)^3 + 2$	44. $y = \frac{1}{2x} - 1$	45. $y = \frac{2}{x^2} + 1$
46. $y = -\sqrt[3]{x^2}$	47. $y = (-2x)^{2/3}$	48. $y = x^2 - 2x $
49. $y = -\sqrt{ x }$	50. $y = 1 + 2\cos(x)$	51. $y = \sin(x) $
52. $y = \frac{1}{4} \tan\left(x - \frac{\pi}{4}\right)$	53. $y = \begin{cases} -x & \text{if } x < 0 \\ e^x - 1 & \text{if } x \ge 0 \end{cases}$	54. $y = 3\ln(x - 2)$

In exercises 55-60, find f^{-1} and it's range of each function

55. $f(x) = \sqrt{10 - 3x}$	56. $f(x) = \frac{4x-1}{2x+3}$	57. $f(x) = e^{x^3}$
58. $f(x) = 2x^3 + 3$	59. $f(x) = \ln(x+3)$	$60. f(x) = \frac{1 + e^x}{1 - e^x}$