

A etiology

- 1. Biological:
- A. Neurotransmitter . L: serotonin
- B. Limbic cyst : lesion
- C. Chromosomal : xyy.
- D. Endocrine : (turmoil change)
 - 1. Androgen so most common occur by men.
 - 2. Premenstrual tension.
- E. Alcohol & Drugs is relate to alcohol intoxicate or withdrawal
- 2. Psychological.
- 1. Developmental factor: the child victim become abusive in future.
- 2. Environmental factors :- weather temperature, overcrowded
- 3. socioeconomic
- 4. Fanatic
- 5. Political

Child abuse

The child injured by abusing physically may cause brain damage & also psychologically as emotion disturbed occur in upper and lower social class.

Clue of child abuse

- 1. Unexplained trauma.
- 2. Delayed in seeking treatment &the parent refuse admit to hospital
- 3. vague somatic symptoms.
- 4. Parental stress.

The people who abused the child: either Alcoholic, depress mother, intellect dull Person.

Sexual abuse

Range from indecent to intercourse, either single meant by stranger or repeated over period by family member or friend of family Than is may be pain, evidence if trauma,.

If between the person that forbidden called (incest) father daughter, brother –sister incest

The father may be by Alcoholic ,personality disorder, psychosis disturb family, mother outside the home ,low I.Q level

Spouse abuse

It is very common in all social classes, most common husband battering wives . 1/4 of homicide occur in family &1/2 of these involve spouse killing spouse .husbands and wives kill each other equally frequently .

Depression, Alcoholism, personality disorder in the cause if abusing partner,

Rape

Sexual intercourse with a female without her consent.

The Definition in restricted

- Some Rapist don't engage in sexy inter course but in other sex like fellatio
 ,or anal sexual
- 2. May be male victim of homosexual.
- 3. May a male victim by female .
- May husband enforced wife to engage in sexily activity.

Incidence if Rape

In USA (1:2000) female.

- -5%f all violence crime.
- -only 1/20 if actual rape repotted to police.
- -most Rapist one married &has
- -year sexual Relationship to their wives.
- -1/2 of case the assault done at the victims' home
- -30-50 % f case the victims is known Rapist –Rapist usually any preset-no semen, or intact hymen.

Causes

- 1. Biological testosterone
- 2. Psychopathic behavior.

Manage

- -Examine the victim if any bruises ,any fracture ,
- -Exam: Depression, anxiety.
- -Treatment the victim, by psychotherapy
- -Treatment the rapist ,rarely come for R

Prevention

- 1. Avoid contact with stranger
- 2. Woman if at a high risk should learn karate

