SOMATOFORM DISORDER

Assistant professor
Dr .Akeel Al sabbagh
Consultant psychiatrist

Somatoform disorder

It is a wide range of syndromes that include the association of physical symptoms (which mimic physical illness) with psychological stress

- 1-Somatization disorder
- 2-Conversion disorder
- 3-Pain disorder
- 4-Hypochondriasis
- 5-Body Dysmorphic disorder

Somatization disorder

History of physical symptoms beginning before age of 30 years over a period of several years and result in causing impairment of social & occupational functioning

Symptoms

Four pain symptoms: pain symptoms (Head, abdominal, Back, joint, chest, rectum, during menstruation, during sexual intercourse

Two gastrointestinal Symptoms: nausea, vomiting, diarrhea

One pseudoneurological: impairment coordination, paralysis, localize weakness, aphonia, urinary retention, Hallucination One sexual symptoms: erectile dysfunction, premature ejaculation, irregular menstruation,

Conversion disorder

It is symptoms affect voluntary motor or sensory function that suggest a neurological or General medical condition

- Usually precede by conflict or stress
- Theses symptoms is not intentionally produced
- After investigation ,the result , not explained a general medical condition

The symptoms cause impairment of social &occupational functioning.

Patient has **2gain**

Primary gain Secondary gain

- -for seeking attention &sympathy
- to avoid responsibilities
- for financial gain
- to release guilt through suffering. Incongruent between mood of the patient and the physical Condition called (la belle indifference)

Pain disorder

It is pain in one or more anatomical sites which the predominant focus of clinical presentation and it is so severe that seeking medical attention.

- -This pain cause impairment of social & occupational functioning.
- -Psychical stress play a role in onset, severity, exacerbation or maintain of the pain.
- -It is not intentionally produce

Hypochondriasis

It's a preoccupation with fear of having or belief of having a disease depend on personals misinterpretation of body symptoms

<u>Diagnostic Guideline</u>

The preoccupant persist even after investigation are normal or after reassurance
The belief is not reach to a delusion.
Causing of social &occupational functioning.
The duration is at least 6 months.

Common region. Head_neck ,,abdomen, chest in order Bodily system: musculoskeletal ,Gastrointestinal, CNS symptoms usually unilateral &mostly on left side.

Bodily dysmorphic disorder

- -It is a preoccupation with the defect in appearance.
- -it is either imagined ,or slightly physical anomaly is present .
- -The person' concern is markedly excessive ,causing impairment of social &occupational functioning
- -It involve usually face ,arm ,genital organ ,It is describe as too small, too lager ,or asymmetrical . It may be associate with social phobia ,so refuse appear in public ,&usually go out after darkness

